

Product Data Sheet



Product Name *3-D Life* Scrambled RGD Peptide

Catalog Number 09-P-003

Description RGD sequences are motifs on extracellular matrix proteins promoting adhesion to cellular integrin receptors. The *3-D Life* Scrambled RGD Peptide contains a scrambled RGD motif and a thiol group on the N-terminus. The thiol group is used to immobilize the peptide on maleimide polymers in *3-D Life* Hydrogels. The immobilized peptide is used in control experiments to determine the specificity of cell adhesion caused by a genuine RGD peptide (e.g. *3-D Life* RGD Peptide).

For instructions, please consult General Protocols GP-1 "Preparation of *3-D Life* Fast Gelling Hydrogels" or GP-2 "Preparation of *3-D Life* Slow Gelling Hydrogels" and the *3-D Life* Hydrogels User Guide on www.cellendes.com.

Quantity 1 μmol

Components

Material	Quantity	Concentration of reactive groups	Storage
● Scrambled RGD Peptide, lyophilized	3x 1 μmol	20 mmol/L*	Lyophilisate and after reconstitution: -20°C to -80°C
○ Water	600 μl	n/a	RT to -80°C

All materials are filter-sterilized.

Reconstitution Scrambled RGD Peptide:

1. Briefly centrifuge vial containing the Scrambled RGD Peptide lyophilisate to make sure that entire product is at the bottom of the reaction tube.
2. Add 48 μl *3-D Life* Water per tube of Scrambled RGD Peptide to dissolve lyophilisate at a concentration of 20 mmol/L thiol groups.
3. Close tube and briefly vortex.
4. Incubate for 5 min.
5. Briefly vortex and centrifuge again.
6. The Scrambled RGD Peptide is now ready for use.

Pept. Sequence Acetyl-Cys-Doa*-Doa-Gly-Arg-Asp-Gly-Ser-Pro-NH₂

*: Doa:8-amino-3,6-dioxaoctanoic acid

Molar Mass 1022 g/mol

Purity >90% (HPLC)

Intended for research use only. Not for use in human therapeutic or diagnostic applications.