

DetachKit / DetachKit 2

PromoCell

Instruction Manual

Product	Size	Catalog Number
DetachKit	30 ml	C-41200
	125 ml	C-41210
	250 ml	C-41220
DetachKit 2	30 ml	C-41202
	125 ml	C-41212
	250 ml	C-41222

Detach-Kit	Detach-Kit 30 C-41200	Detach-Kit 125 C-41210	Detach-Kit 250 C-41220
Contains	HEPES-BSS C-40000 Trypsin/EDTA C-41000 TNS C-41100	C-40010 C-41010 C-41110	C-40020 C-41020 C-41120

Detach-Kit 2	Detach-Kit 2-30 C-41202	Detach-Kit 2-125 C-41212	Detach-Kit 2-250 C-41222
Contains	HEPES-BSS C-40000 Trypsin/EDTA 2 C-41002 TNS C-41100	C-40010 C-41012 C-41110	C-40020 C-41022 C-41120

Product Description

PromoCell DetachKits were designed for the safe and efficient detachment of primary human cells in routine subculturing. DetachKit 2 is especially valuable for more sensitive cell types.

Each DetachKit consists of three components: HEPES-BSS (HEPES-buffered Balanced Salt Solution), Trypsin/EDTA Solution and TNS (Trypsin Neutralization Solution).

The DetachKit comes with Trypsin/EDTA (0.04%/0.03%) whereas the DetachKit 2 contains a lower concentration of Trypsin/EDTA (0.025%/0.01%). HEPES-BSS contains 30 mM HEPES,

D-Glucose, NaCl, KCl, Na-Phosphate and Phenol Red. TNS contains 0.05% Trypsin Inhibitor from soybean and 0.1% Bovine Serum Albumin.

Storage and Stability

Store at -20°C in the dark immediately after arrival. Refrigerate thawed solution at 4 to 8°C. If stored properly, the product is stable until the expiry date stated on the label.

Quality Control

All lots of PromoCells DetachKits are subjected to comprehensive quality

control tests. Each lot is routinely tested for use with primary human cells, the absence of cytotoxicity, and physical parameters such as osmolality and pH level. Approved in-house lots are used as a reference.

In addition, all lots have been tested for the absence of microbial contaminants (fungi, bacteria).

Intended Use

The products are for *in vitro* research use only and not for diagnostic or therapeutic procedures. For safety precautions please see appropriate MSDS.

Subcultivation Protocol


1. Prepare the reagents and wash the cells

Place the PromoCell DetachKit at room temperature for at least 30 minutes to adjust the temperature of the reagents. Carefully aspirate the medium from the culture vessel. Add 100 μ l HEPES BSS Solution per cm^2 of vessel surface to wash the cells and agitate the vessel carefully for 15 seconds.



2. Detach the cells

Carefully aspirate the HEPES BSS from the culture vessel. Add 100 μ l Trypsin/EDTA Solution per cm^2 of vessel surface. Note: We recommend detaching the cells at room temperature. Close the vessel and examine the cells under a microscope. When the cells start to detach, gently tap the side of the vessel to loosen the remaining cells.



3. Neutralize the trypsin and harvest the cells

Add 100 μ l Trypsin Neutralization Solution per cm^2 of vessel surface and gently agitate. Carefully aspirate the cell suspension and transfer it to a centrifugation tube. Spin down the cells for 3 minutes at 220 x g.



4. Incubate the cells

Discard the supernatant (step 1), add 1 ml of the appropriate PromoCell Cell Growth Medium (step 2), and resuspend the cells by carefully pipetting up and down. Plate the cells according to the recommended seeding density in new cell culture vessels containing PromoCell Cell Growth Medium pre-warmed to 37°C. Place the vessels in an incubator (37°C, 5% CO_2).



PromoCell GmbH

Sickingenstr. 63/65
69126 Heidelberg
Germany

Email: info@promocell.com
www.promocell.com

USA/Canada

Phone: 1 – 866 – 251 – 2860 (toll free)
Fax: 1 – 866 – 827 – 9219 (toll free)

Deutschland

Telefon: 0800 – 776 66 23 (gebührenfrei)
Fax: 0800 – 100 83 06 (gebührenfrei)

France

Téléphone: 0800 90 93 32 (ligne verte)
Téléfax: 0800 90 27 36 (ligne verte)

United Kingdom

Phone: 0800 – 96 03 33 (toll free)
Fax: 0800 – 169 85 54 (toll free)

Other Countries

Phone: +49 6221 – 649 34 0
Fax: +49 6221 – 649 34 40