



Multiplex IHC: Making Discoveries Multicolor

Complexity of the Tumor Microenvironment

The tumor microenvironment (TME) is a complex mass of malignant and nonmalignant cells, signaling molecules, extracellular matrix, and blood vessels. Inhibition of immune checkpoints and co-inhibitory molecules such as CTLA-4 and PD-1 to immunomodulate the T-cell response within the TME is a promising cancer therapy. Multiplex immunohistochemistry (mIHC) enables the tracking of multiple markers within the TME, predicting therapeutic response and highlighting new therapeutic targets

TME Expression Profiles

UNDERSTANDING OF THE INTERACTIONS BETWEEN MALIGNANT AND NONMALIGNANT CELLS.

TME expression profiles guide understanding of the interactions between malignant and nonmalignant cells.

T cells: CD3, CD4, PD-1, CTLA-4, FoxP3, CD4, granzyme B, granzyme A, CD25, CD39, CD73, CD103

Extracellular matrix: collagen, fibronectin, laminin

Macrophages: CD68, HLA-DR, CD14, CD11b, CD163, CX3CR1


B cells: CD19, CD20, CD40, CD80, CD86, CD69

CUSTOM PUBLISHING FROM
TheScientist
EXPLORING LIFE, INSPIRING INNOVATION

Traditional One-Color IHC


Traditional, one-color IHC can not differentiate between multiple cell types and targets in tumors. To immunostain more than two targets, serial sections or serial staining procedures must be used.

One-Color Process



Traditional mIHC


Without Amplification mIHC Process



Tyramide Signal Amplification mIHC

SEEING LIFE IN MULTICOLOR

With Amplification mIHC Process



mIHC allows multiple targets to be labeled. Traditional mIHC uses primary antibodies from different species. Tyramide signal amplification mIHC use multiple rounds of staining and antibody removal.

Multiplex IHC for Immuno-oncology: Seeing Life in Multicolor

Sequential staining of different antigens with:

- Primary antibody
- HRP conjugated secondary antibody
- Tyramide-conjugated fluorochromes

Tyramide labels adjacent to the antigens and binds covalently. Primary and Secondary antibodies are removed by microwaving after each round of staining.

ADVANTAGES OF MULTIPLEX IHC

1. Multiplex IHC allows for visualization of multiple targets within a single tissue section, critical for limited samples.

2. Tissue architecture is preserved allowing for spatial information and co-expression within the TME, unlike alternative multiplex approaches such as NGS, PCR, mass spectrometry, etc.

3. Fluorophore detection systems offer major advantages over chromogenic detection:

- Fluorophores have a wider dynamic range and larger linear range than chromogenic substrates, tyramide-based multiplexing enhances fluorescence signal enabling detection of low-level binding sites.
- DAPI (DNA/nuclear counterstain) is superior to hematoxylin, which can be obscured by other targets with chromogenic staining.
- Fluorescence signals can be overlaid and seen as single or multi-channel, allowing for intensity measurements for each target.