

DESCRIPTION

Source *Spodoptera frugiperda*, Sf 21 (baculovirus)-derived
Thr32-His266
Accession # O94907

N-terminal Sequence Analysis Thr32

Predicted Molecular Mass 25.8 kDa

SPECIFICATIONS

SDS-PAGE 33-40 kDa, reducing conditions

Activity Measured by its ability to inhibit Wnt-3a-induced alkaline phosphatase production by MC3T3-E1 mouse preosteoblast cells. The ED₅₀ for this effect is typically 0.03-0.15 µg/mL in the presence of 10 ng/mL of rmWnt-3a.

Endotoxin Level <1.0 EU per 1 µg of the protein by the LAL method.

Purity >95%, by SDS-PAGE under reducing conditions and visualized by silver stain.

Formulation Lyophilized from a 0.2 µm filtered solution in PBS. See Certificate of Analysis for details.

PREPARATION AND STORAGE

Reconstitution Reconstitute at 100 µg/mL in PBS.

Shipping The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Dickkopf related protein 1 (Dkk-1) is the founding member of the Dickkopf family of proteins that includes Dkk-1, -2, -3, -4, and a related protein, Soggy (1, 2). Dkk proteins are secreted proteins that contain two conserved cysteine-rich domains separated by a linker region. Each domain contains ten cysteine residues (1 - 3). Mature human Dkk-1 is a 40 kDa glycosylated protein that shares 86%, 87%, 90% and 91% aa sequence identity with mouse, rat, rabbit and bovine Dkk-1, respectively. It also shares 42% and 36% aa identity with human Dkk-2 and Dkk-4, respectively. Dkk-1 and Dkk-4 are well documented antagonists of the canonical Wnt signaling pathway (1, 2). This pathway is activated by Wnt engagement of a receptor complex composed of the Frizzled proteins and one of two low-density lipoprotein receptor-related proteins, LRP5 or LRP6 (4). Dkk-1 antagonizes Wnt by forming ternary complexes of LRP5/6 with Kremen1 or Kremen2 (4, 5). Dkk-1/LRP6/Krm2 complex internalization has been shown to downregulate Wnt signaling (4, 5). Dkk-1 is expressed throughout development and antagonizes Wnt-7a during limb development (6, 7). Other sites of expression include developing neurons, hair follicles and the retina of the eye (8, 9). The balance between Wnt signaling and Dkk-1 inhibition is critical for bone formation and homeostasis (10). Insufficient or excess Dkk-1 activity in bone results in increased or decreased bone density, respectively (8, 11). In adults, Dkk-1 is expressed in osteoblasts and osteocytes, and neurons. Cerebral ischemia induces Dkk-1 expression, which contributes to neuronal cell death (12).

References:

1. Krupnik, V.E. *et al.* (1999) *Gene* **238**:301.
2. Niehrs, C. (2006) *Oncogene* **25**:7469.
3. Bullock, C.M. *et al.* (2004) *Mol. Pharmacol.* **65**:582.
4. Mao, B. *et al.* (2001) *Nature* **411**:321.
5. Mao, B. *et al.* (2002) *Nature* **417**:664.
6. Kemp, C. *et al.* (2005) *Dev. Dyn.* **233**:1064.
7. Adamska, M. *et al.* (2004) *Dev. Biol.* **272**:134.
8. Li, J. *et al.* (2006) *Bone* **36**:754.
9. Verani, R. *et al.* (2006) *J. Neurochem.* **101**:242.
10. Pinzone, J.J. *et al.* (2009) *Blood* **113**:517.
11. Morvan, F. *et al.* (2006) *J. Bone Miner. Res.* **21**:934.
12. Cappuccio, I. *et al.* (2005) *J. Neurosci.* **25**:2647.