

## DESCRIPTION


<b>Species Reactivity</b>	Human/Mouse
<b>Specificity</b>	Detects endogenous human and mouse NCOA3 in Western blots.
<b>Source</b>	Polyclonal Goat IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human NCOA3 Met673-Val856 Accession # Q9Y6Q9
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	0.1 µg/mL	See Below

## DATA



## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.2 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<p><b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b></p> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month from date of receipt, 2 to 8 °C, reconstituted.</li> <li>● 6 months from date of receipt, -20 to -70 °C, reconstituted.</li> </ul>

## BACKGROUND

Nuclear Receptor Coactivator 3 (NCOA3) functions in the multisubunit coactivator complex. Multiple nuclear hormone receptors are able to bind to NCOA3 in a hormone dependent manner. NCOA3 contains intrinsic histone acetyltransferase activity, but is also able to recruit other histone acetyltransferases (CBP, PCAF) and methyltransferases (PRMT1, CARM1) to active promoters. NCOA3 null mice exhibit hematological, adipogenesis, and endotoxin impairments. mRNA of NCOA3 is often amplified in several cancers including breast and ovarian.