

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human FABP8 in direct ELISAs and Western blots. In direct ELISAs, less than 2% cross-reactivity with recombinant human (rh) FABP2, 3, 4, 5, 6, 7 and 9 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human FABP8 Ser2-Val132 Accession # P02689
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	See Below
Immunohistochemistry	5-15 µg/mL	See Below


DATA

Immunohistochemistry




FABP8 in Human Spinal Cord. FABP8 was detected in immersion fixed paraffin-embedded sections of human spinal cord using 15 µg/mL Human FABP8 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF5866) overnight at 4 °C. Tissue was stained with the Anti-Sheep HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS019) and counterstained with hematoxylin (blue). View our protocol for [Chromogenic IHC Staining of Paraffin-embedded Tissue Sections](#).

Immunohistochemistry



FABP8 in Human Brain. FABP8 was detected in immersion fixed paraffin-embedded sections of human brain (cortex) using Human FABP8 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF5866) at 15 µg/mL overnight at 4 °C. Tissue was stained using the Anti-Sheep HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS019) and counterstained with hematoxylin (blue). View our protocol for [Chromogenic IHC Staining of Paraffin-embedded Tissue Sections](#).

Western Blot



Detection of Human FABP8 by Western Blot. Western blot shows lysates of human hippocampal and rat spinal cord tissue. PVDF membrane was probed with 1 µg/mL of Human FABP8 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF5866) followed by HRP-conjugated Anti-Sheep IgG Secondary Antibody (Catalog # HAF016). A specific band was detected for FABP8 at approximately 15 kDa (as indicated). This experiment was conducted under reducing conditions and using [Immunoblot Buffer Group 7](#).

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month from date of receipt, 2 to 8 °C, reconstituted. ● 6 months from date of receipt, -20 to -70 °C, reconstituted.

BACKGROUND

FABP8 (fatty acid binding protein-8; also M [myelin]-FABP, P2 and PMP2) is a 15 kDa (predicted) member of the fatty acid binding protein family, calycin superfamily of molecules. It is found in Schwann cells, presumably on the cytoplasmic face of the plasma membrane where it may contribute to fatty acid transport across myelin. Functionally, FABP8 has a high affinity for U-shaped fatty acids such as oleic and palmitic acid. Human FABP8 is 132 amino acids (aa) in length and exhibits two layers of anti-parallel β -strands that envelope a hydrophobic pocket for lipid binding. Arg107 plus Arg127-Ile128-Tyr129 participate in fatty acid binding. Full length human FABP8 shares 87% and 95% aa identity with mouse and rabbit FABP8, respectively.