

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Chitobiase/CTBS in direct ELISAs and Western blots.
Source	Monoclonal Mouse IgG ₁ Clone # 486725
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Chitobiase/CTBS Thr39-Arg385 Accession # Q01459
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	Recombinant Human Chitobiase/CTBS
Immunoprecipitation	25 µg/mL	Conditioned cell culture medium spiked with Recombinant Human Chitobiase/CTBS, see our available Western blot detection antibodies

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Di-N-acetyl-chitobiase (CTBS) is a lysosomal glycosidase that is involved in the degradation of asparagine-linked glycoproteins. The amino acid sequence of human CTBS shares 99%, 86%, 82%, and 77% identity with that of chimpanzee, mouse, rat, and canine CTBS, respectively.