

Anti-Pf-FNR (Ferredoxin--NADP reductase, P. falciparum) antibody, rabbit polyclonal

81-009 400 μg

Storage: Ship at 4°C and store at -20°C. Do not freeze.

Immunogen: Purified recombinant *P. falciparum* Ferredoxin-NADP reductase (full-size, no-tag attached) expressed in *E. coli.*

Reactivity: FNR protein of *Plasmodium falciparum*. Cross-reacts also with plant FNR isoproteins.

Applications:

- 1. Western blotting (1/500-1/2,000 dilution). Extract for western blotting should be made from apicoplast fraction of *P. falciparum*.
- 2. ELISA (assay dependent)

Purity: Protein A purified IgG

Form: 4 mg/ml in PBS, 50% glycerol. Filter sterilized. Azide and carrier free.

Background: Fd:NADPH oxidoreducatase (FNR) plays a key role in regulating the relative amounts of cyclic and non-cyclic electron flow to meet the demands of the plant for ATP and reducing power. The human malaria parasite (Plasmodium falciparum) possesses a plastid-derived organelle called the apicoplast, which is believed to employ metabolisms crucial for the parasite's survival.

Sucellular location: Apicoplast (plastid-like organelle)

Data Link: UniProKB <u>C6KT68</u> (FENR_PLAF7)




Fig.1 Western blot of Ferredoxin--NADP reductase of *P. falciparum* with anti-*Pf* FNR antibody.

Anti-Pf FNR antibody was used at 1/1,000 dilution. Second antibody (goat anti-rabbit IgG antibody HRP-conjugated, ab97051) was used at 1/10,000 dilution.

Sample; 1 µl of 40 µM recombinant pfFNR

Molecular mass indicated from the gene is 43,8 kDa. However, transit peptide consisting of N-terminal 18 amino acids is removed in the mature form.






Fig.2 The anti-Pf-FNR antibody reacts also with plant FNR proteins in western blotting.

- 1. Extract of Arabidopsis leaf (10 ug)
- 2. Extract of Maize leaf (10 ug)

The antibody was used at 1/1,000 dilution Astrerisk indicates a nonspecific band.

Reference: Recombinant *Pf* FNR is described in the following publication.

1.Kimata-Ariga Y et al. Cloning and characterization of ferredoxin and ferredoxin-NADP+ reductase from human malaria parasite. <u>J Biochem.</u> 2007 Mar;141(3):421-8 PMID:17251200. **WB, IF;** *P. falciparum.*