DEC1 IHC Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP 003661.1

Catalog No. IHC-00448-T Gene ID 8553



APPLICATIONS IHC-P, IF

REACTIVITY TESTED Human. Other species have not been tested.

PRESUMED REACTIVITY Based on 100% sequence identity, this antibody is predicted to react with Orangutan, Rhesus Monkey,

Gorilla, Chimpanzee, Crab-eating macaque and Northern white-cheeked gibbon.

ISOTYPE IgG

AMOUNT 10 μl

STORAGE/SHELF LIFE 2 - 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-buffered Saline containing 0.1% BSA containing 0.09% Sodium Azide

ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to DEC1 immobilized on solid support.

The epitope recognized by IHC-00448-T maps to a region between residues 350 and the C-terminus (residue 412) of human Differentially Expressed in Chondrocytes (Basic Helix-Look-Helix Domain

Containing, Class B, 2) using the numbering given in entry NP_003661.1 (GeneID 8553).

Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Immunohistochemistry 1:100 - 1:500 Immunofluorescence 1:50 - 1:500

APPLICATION NOTES Epitope exposure is recommended.

Epitope exposure with tris-EDTA pH9. 0 buffer will enhance staining.


Likely to work with frozen sections.

In some cases, the antibody may be diluted further than indicated.

IHC HUMAN CONTROLS Breast Carcinoma, Ovarian Carcinoma, Prostate Carcinoma

ADDITIONAL INFO http://www.bethyl.com/product/IHC-00448-T

Use the link above to view SDS, a current list of citations, and other product specific information.



Detection of Human Dec1 by Immunohistochemistry. *Sample:* FFPE section of human breast carcinoma. *Antibody:* Affinity purified rabbit anti-Dec1 (Cat. No. IHC-00448-T) used at a dilution of 1:250. *Detection:* DAB staining using Immunohistochemistry Accessory Kit (Cat. No. IHC-101).