

MDM2 Polyclonal Antibody, Cy5 Conjugated

DATASHEET

Host: Rabbit

Target Protein: MDM2

Immunogen 440-491/491

Range:

Clonality: Polyclonal

Isotype: IgG

Entrez Gene: [4193](#)

Swiss Prot: [Q00987](#)

Source: KLH conjugated synthetic peptide derived from human MDM2

Purification: Purified by Protein A.

Storage: Aqueous buffered solution containing 100ug/ml BSA, 50% glycerol and 0.09% sodium azide. Store at 4°C for 12 months.

Background: E3 ubiquitin-protein ligase that mediates ubiquitination of p53/TP53, leading to its degradation by the proteasome. Inhibits p53/TP53- and p73/TP73-mediated cell cycle arrest and apoptosis by binding its transcriptional activation domain. Also acts as a ubiquitin ligase E3 toward itself and ARRB1. Permits the nuclear export of p53/TP53. Promotes proteasome-dependent ubiquitin-independent degradation of retinoblastoma RB1 protein. Inhibits DAXX-mediated apoptosis by inducing its ubiquitination and degradation. Component of the TRIM28/KAP1-MDM2-p53/TP53 complex involved in stabilizing p53/TP53. Also component of the TRIM28/KAP1-ERBB4-MDM2 complex which links growth factor and DNA damage response pathways. Mediates ubiquitination and subsequent proteasome degradation of DYRK2 in nucleus. Ubiquitinates IGF1R and SNAI1 and promotes them to proteasomal degradation.

Conjugation: Cy5

Excitation/ Emission: 625,650nm/670nm

Size: 100ul

Concentration: 1ug/ul

Applications: IF(IHC-P)(1:50-200)

Cross Reactive Species: Human
 Mouse
 Rat

For research use only. Not intended for diagnostic or therapeutic use.