


Rabbit Anti-ATG16L Polyclonal Antibody, FITC Conjugated



www.biossusa.com support@biossusa.com 800.501.7654 [DOMESTIC] +1.781.569.5821 [INTERNATIONAL]

— D A	T ASHEET	
Host:	Rabbit	Conjugation: FITC
Target Protein:	ATG16L	Excitation/ 494nm/518nm
•	530-580/607	Emission:
Range: Clonality:	Polyclonal	Size: 100ul
Isotype:	IgG	Concentration: 1ug/ul
Entrez Gene:	55054	Applications: IF(IHC-P)(1:50-200)
Swiss Prot:	Q676U5	Cross Reactive Human Species: Mouse
Source:	KLH conjugated synthetic peptide derived from human ATG16A	Rat
Purification:	Purified by Protein A.	Caution: For research use only. Not for human or animal
Storage:	Aqueous buffered solution containing 100ug/ml BSA, 50% glycerol and 0.09% sodium azide. Store at 4°C for 12 months.	therapeutic or diagnostic use.
Background:	Plays an essential role in autophagy: interacts with ATG12-ATG5 to mediate the conjugation of phosphatidylethanolamine (PE) to LC3 (MAP1LC3A, MAP1LC3B or MAP1LC3C), to produce a membrane-bound activated form of LC3 named LC3-II. Thereby, controls the elongation of the nascent autophagosomal membrane.	
— V — A	L I D A T I O N	
I M A G E S		



Formalin-fixed and paraffin embedded mouse spleen labeled with Anti-ATG16L Polyclonal Antibody, Unconjugated (bs-4007R) at 1:200 followed by conjugation to the secondary antibody and DAB staining