



11-143-C100

Monoclonal Antibody to CD137 / 4-1BB Purified Antibody (0.1 mg)

Clone:	4B4-1
Isotype:	Mouse IgG1
Specificity:	The mouse monoclonal antibody 4B4-1 recognizes CD137, an approximately 40 kDa type I transmembrane protein of the TNFR family expressed mainly on activated T cells. HLDA VI; WS Code C-7
Regulatory Status:	RUO
Immunogen:	Recombinant human CD137 ectodomain
Species Reactivity:	Human, Non-Human Primates
Application:	Flow Cytometry Immunoprecipitation ELISA Functional Application inhibition of cytokine production
Purity:	> 95% (by SDS-PAGE)
Purification:	Purified by protein-A affinity chromatography
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD137, also known as TNFRSF9 or 4-1BB, is an inducible costimulatory molecule expressed mainly on activated T cells. Its ligand, known as 4-1BBL, is expressed on activated macrophages, mature B cells, hematopoietic stem cells, and myeloid progenitor cells. CD137 signaling leads to maintaining the survival of activated T cells and CD8+ memory T cells, and clonal expansion of T cells, but also to suppressing myelopoiesis and dendritic cell development. Triggered CD137 induces a cytokine release profile regulating peripheral monocyte survival. Soluble forms of CD137 may provide negative control mechanism for some immune responses.

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

- References:**
- *Chan SL, Voskens CJ, Lin W, Schindler DG, Azimzadeh A, Wang LX, Taylor RJ, Strome SE, Schulze DH: Epitope mapping of a chimeric CD137 mAb: a necessary step for assessing the biologic relevance of non-human primate models. *J Mol Recognit.* 2009 May-Jun;22(3):242-9. doi: 10.1002/jmr.937.
 - *Yi L, Zhao Y, Wang X, Dai M, Hellström KE, Hellström I, Zhang H: Human and mouse CD137 have predominantly different binding CRDs to their respective ligands. *PLoS One.* 2014 Jan 21;9(1):e86337.
 - *Fernández Do Porto DA, Jurado JO, Pasquinelli V, Alvarez IB, Aspera RH, Musella RM, García VE: CD137 differentially regulates innate and adaptive immunity against *Mycobacterium tuberculosis*. *Immunol Cell Biol.* 2012 Apr;90(4):449-56.
 - *Bellarosa D, Bressan A, Bigioni M, Parlani M, Maggi CA, Binaschi M: SAHA/Vorinostat induces the expression of the CD137 receptor/ligand system and enhances apoptosis mediated by soluble CD137 receptor in a human breast cancer cell line. *Int J Oncol.* 2012 Oct;41(4):1486-94.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz