



11-760-C100

Monoclonal Antibody to CD116 Purified Antibody (0.1 mg)

Clone:	4H1
Isotype:	Mouse IgG1
Specificity:	The mouse monoclonal antibody 4H1 recognizes human CD116, the GM-CSF receptor alpha subunit (approx. 80 kDa) expressed e.g. by neutrophils, eosinophils, monocytes and macrophages.
Regulatory Status:	RUO
Immunogen:	CD116-transfected COS cells
Species Reactivity:	Human
Application:	Flow Cytometry Immunoprecipitation Western Blotting Immunohistochemistry (frozen sections) Mass Cytometry
Purity:	> 95% (by SDS-PAGE)
Purification:	Purified by protein-A affinity chromatography
Concentration:	1 mg/ml
Storage Buffer:	Tris buffered saline (TBS) with 15 mM sodium azide, approx. pH 8.0
Storage / Stability:	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD116 (GM-CSF R alpha) is the low affinity receptor for granulocyte-macrophage colony-stimulating factor (GM-CSF). CD116 heterodimerizes with CD131, the common beta chain subunit shared with IL-3 and IL5 receptors, to form the high affinity GM-CSF receptor. CD116 is expressed by myeloid cells including macrophages, neutrophils, eosinophils, dendritic cells, and their precursors, as well as on endothelial cells. It is being used as a specific marker of myeloid leukemias.

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

- References:**
- Stomski FC, Woodcock JM, Zacharakis B, Bagley CJ, Sun Q, Lopez AF: Identification of a Cys motif in the common beta chain of the interleukin 3, granulocyte-macrophage colony-stimulating factor, and interleukin 5 receptors essential for disulfide-linked receptor heterodimerization and activation of all three receptors. *J Biol Chem.* 1998 Jan 9;273(2):1192-9.
 - *Huntington ND, Legrand N, Alves NL, Jaron B, Weijer K, Plet A, Corcuff E, Mortier E, Jacques Y, Spits H, Di Santo JP: IL-15 trans-presentation promotes human NK cell development and differentiation in vivo. *J Exp Med.* 2009 Jan 16;206(1):25-34
 - *Koba C, Haruta M, Matsunaga Y, Matsumura K, Haga E, Sasaki Y, Ikeda T, Takamatsu K, Nishimura Y, Senju S: Therapeutic effect of human iPS-cell-derived myeloid cells expressing IFN- γ ; against peritoneally disseminated cancer in xenograft models. *PLoS One.* 2013 Jun 24;8(6):e67567
 - *Schwarzmaier D, Foell D, Weinhage T, Varga G, Däbritz J: Peripheral monocyte functions and activation in patients with quiescent Crohn's disease. *PLoS One.* 2013 Apr 26;8(4):e62761.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz