



11-777-C025

## Monoclonal Antibody to CD112 Purified Antibody (0.025 mg)

<b>Clone:</b>	R2.525
<b>Isotype:</b>	Mouse IgG1
<b>Specificity:</b>	The mouse monoclonal antibody R2.525 recognizes CD112, a type I transmembrane glycoprotein expressed by myelomonocytic and megakaryocytic cells, and by CD34+ hematopoietic progenitors.
<b>Regulatory Status:</b>	RUO
<b>Immunogen:</b>	NIH/3T3 cells transfected with human Nectin-2
<b>Species Reactivity:</b>	Human
<b>Application:</b>	Flow Cytometry Immunoprecipitation Immunohistochemistry (frozen sections)
<b>Purity:</b>	> 95% (by SDS-PAGE)
<b>Purification:</b>	Purified by protein-A affinity chromatography
<b>Concentration:</b>	1 mg/ml
<b>Storage Buffer:</b>	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
<b>Storage / Stability:</b>	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label
<b>Background:</b>	CD112, also known as nectin-2, is a transmembrane glycoprotein involved in organization of adherens junctions. It also serves as a target molecule for entry of certain strains of herpes simplex virus (HSV) and pseudorabies virus (PRV). It is homologous to CD155, which serves as a target molecule for polio virus. CD112 seems to play a role in neural tube formation, with N-cadherin. Inside the cell, CD112 is connected with actin cytoskeleton through afadin. Variations in the CD112 gene have been associated with differences in the severity of multiple sclerosis. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.

**For laboratory research only, not for drug, diagnostic or other use.**



**Antibodies**

**References:**

\*Enqvist M, Nilsson G, Hammarfjord O, Wallin RP, Björkström NK, Björnstedt M, Hjerpe A, Ljunggren HG, Dobra K, Malmberg KJ, Carlsten M: Selenite induces posttranscriptional blockade of HLA-E expression and sensitizes tumor cells to CD94/NKG2A-positive NK cells. *J Immunol.* 2011 Oct 1;187(7):3546-54.

Nielsen N, &#216;dum N, Urs&#248; B, Lanier LL, Spee P: Cytotoxicity of CD56(bright) NK cells towards autologous activated CD4+ T cells is mediated through NKG2D, LFA-1 and TRAIL and dampened via CD94/NKG2A. *PLoS One.* 2012;7(2):e31959.

\*Carlsten M, Björkström NK, Norell H, Bryceson Y, van Hall T, Baumann BC, Hanson M, Schedvins K, Kiessling R, Ljunggren HG, Malmberg KJ: DNAX accessory molecule-1 mediated recognition of freshly isolated ovarian carcinoma by resting natural killer cells. *Cancer Res.* 2007 Feb 1;67(3):1317-25.

\*Hou S, Ge K, Zheng X, Wei H, Sun R, Tian Z: CD226 protein is involved in immune synapse formation and triggers Natural Killer (NK) cell activation via its first extracellular domain. *J Biol Chem.* 2014 Mar 7;289(10):6969-77.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at [www.exbio.cz](http://www.exbio.cz).

**For laboratory research only, not for drug, diagnostic or other use.**

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic

Tel: +420 261 090 666 | Fax: +420 261 090 660 | [orders@exbio.cz](mailto:orders@exbio.cz) | [www.exbio.cz](http://www.exbio.cz)