



1P-784-T025

Monoclonal Antibody to CD235a Phycoerythrin (PE) conjugated (25 tests)

| | |
|-----------------------------|--|
| Clone: | JC159 |
| Isotype: | Mouse IgG1 |
| Specificity: | The mouse monoclonal antibody JC159 recognizes an epitope between amino acids 27 and 40 of the extracellular portion of CD235a (glycophorin A), a sialoglycoprotein expressed on early erythroblasts, late erythroblasts, erythroblasts, mature erythrocytes and the cells of erythroid cell lines K562 and HEL. The antibody does not react with glycophorin B. |
| Regulatory Status: | RUO |
| Immunogen: | Membrane preparation from splenic hairy cell leukemia |
| Species Reactivity: | Human, Rat |
| Preparation: | The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary. |
| Storage Buffer: | The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide. |
| Storage / Stability: | Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label. |
| Usage: | The reagent is designed for Flow Cytometry analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.25 ml) is sufficient for 25 tests. |
| Expiration: | See vial label |
| Lot Number: | See vial label |
| Background: | CD235a (Glycophorin A, GPA) is a transmembrane sialoglycoprotein expressed on erythrocytes and their precursors. Similarly to glycophorin B (GPB), these molecules provide the cells with a large mucin-like surface, which minimalizes aggregation between erythrocytes in the circulation. GPA is the carrier of blood group M and N specificities, while GPB accounts for S, s and U specificities. CD235a is a receptor of Hsa, an Streptococcus adhesin. |
| References: | *Maijenburg MW, Kleijer M, Vermeul K, Mul EP, van Alphen FP, van der Schoot CE, Voermans C: The composition of the mesenchymal stromal cell compartment in human bone marrow changes during development and aging. <i>Haematologica</i> . 2012 Feb;97(2):179-83. *Beck Z, Jagodzinski LL, Eller MA, Thelian D, Matyas GR, Kunz AN, Alving CR: Platelets and erythrocyte-bound platelets bind infectious HIV-1 in plasma of chronically infected patients. <i>PLoS One</i> . 2013 Nov 25;8(11):e81002. *Yamauchi T, Takenaka K, Urata S, Shima T, Kikushige Y, Tokuyama T, Iwamoto C, Nishihara M, Iwasaki H, Miyamoto T, Honma N, Nakao M, Matozaki T, Akashi K: Polymorphic Sirpa is the genetic determinant for NOD-based mouse lines to achieve efficient human cell engraftment. <i>Blood</i> . 2013 Feb 21;121(8):1316-25. *Alijotas-Reig J, Palacio-Garcia C, Llurba E, Vilardell-Tarres M: Cell-derived microparticles and vascular pregnancy complications: a systematic and comprehensive review. <i>Fertil Steril</i> . 2013 Feb;99(2):441-9. |

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz