

Product Datasheet

Quinacrine Inhibitor NBP2-29385-25g

Unit Size: 25 g

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-29385

Updated 2/18/2018 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-29385



NBP2-29385-25g


Quinacrine Inhibitor

Product Information	
Unit Size	25 g
Concentration	Concentration is not relevant for this product. Please see the protocols for proper use of this product.
Storage	Store at 4C. Do not freeze.
Buffer	Form : Yellow Solid (Powder) Solubility:Deionized water
Product Description	
Immunogen	CAS Number 69-05-6 Empirical Formula (Hill Notation) C ₂₃ H ₃₀ CIN ₃ O. 2HCl Molecular Weight 472.88
Details of Functionality	Endosomal toll-like receptor inhibitor (antagonist); Inhibitor of endosomal acidification on which functional activity of endosomal TLRs (particularly TLR9 and TLR3) is dependent.
Inhibitor Family	TLR
Product Application Details	
Application Notes	Assay>=90%




Images


Quinacrine Inhibitor [NBP2-29385]



Quinacrine Inhibitor [NBP2-29385] - Evaluation of inhibitory activity of quinacrine. TLR9/NF- κ B/SEAPorter™ HEK 293 and TLR2/NF- κ B/SEAPorter™ HEK 293 cells were plated in 96-well plates at 5×10^4 cells/well for 16 h. Cells were preincubated with different concentrations of quinacrine (0.05, 0.1, 0.5, 1, 5 and 10 μ M) for 30 min. Cells were then stimulated with 10 μ g/ml CpG [A] or 20 ng/ml MALP-2 [B] for 24 h. Secreted alkaline phosphatase (SEAP) was analyzed using SEAPorter™ Assay Kit



Quinacrine Inhibitor [NBP2-29385] - TLR9/NF- κ B/SEAPorter™ HEK 293 cells were plated in 96-well plates at 5×10^4 cells/well for 16 h. Cells were preincubated with various concentrations of quinacrine for 30 min, followed by stimulation with 10 μ g/ml CpG for 24 h. SEAP was analyzed using SEAPorter™ Assay Kit. Dose-responsive percent inhibition of each sample well was calculated to yield the quinacrine IC₅₀ value.



Procedures

Product Handling Protocol (NBP2-29385)

1. To make 100 mM stock solution, dissolve 200 mg quinacrine in 4.2 ml water by gentle vortex.
2. Filter sterilize through a 0.22 um filter.
3. Store at 4oC (Note: Quinacrine solution is light sensitive.).
4. For TLR signaling inhibition study, perform a pilot inhibitory test with the different concentrations of quinacrine ranging from 1 to 10 mM to optimize your experiments.



Novus Biologicals USA

8100 Southpark Way, A-8
Littleton, CO 80120
USA

Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
novus@novusbio.com

Novus Biologicals Canada

461 North Service Road West, Unit B37
Oakville, ON L6M 2V5
Canada

Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada@novusbio.com

Novus Biologicals Europe

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: technical@novusbio.com
Orders: orders@novusbio.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Inhibitors are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-29385

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

