

DATA SHEET for dBcrp, ABCG2, ProVesicles

dBcrp Sf9-membrane derived vesicles for uptake

Cat. #PV31006; Lot #PN-V-1001


Contents: 500 μ L @ 5 mg/mL total protein (determined by BCA protein assay) in MRP/BCRP Resuspension Buffer*

*MRP/BCRP Resuspension Buffer: 50 mM Tris-HCl, pH 7.0; 50 mM mannitol; 2 mM EGTA; 2 mM DTT; 8 μ g/mL aprotinin and 10 μ g/mL leupeptin.

Store at -70 to -80C upon receipt. Aliquot to smaller working volumes to minimize freeze-thawing cycles.

Dog breast cancer resistance protein (dBcrp) inside-out vesicles are prepared from Sf9 insect cells infected with baculovirus to overexpress Bcrp. ProNovus dBcrp ProVesicles should be used to investigate drug interactions with dBcrp *in vitro*.

Representative data showing time dependence, concentration dependence and competitive inhibition of uptake at dBcrp when using Lucifer Yellow (LY), a fluorescent substrate:



ATP-dependent Lucifer Yellow (10 μ M) uptake: 500 pmol/mg/min

Vesicle Uptake Assay Protocol:

1. Incubate a 95 μ L reaction containing 50 μ g vesicles and LY in MRP/BCRP Uptake Buffer (10 mM Tris-HCl, pH 7.4; 250 mM sucrose; 10 mM $MgCl_2$) for 5 min at 37C.
2. Uptake was started by adding MgATP (or AMP) at a final of 5 mM and incubating for 5 min at 37C.
3. Reaction was terminated by transferring vesicle samples to a filter plate and washing the filter plate 6 x with ice cold washing buffer (40 mM MOPS-Tris, pH 7.0; 70 mM KCl).
4. Filter plate was dried and fluorescent counts measured with a spectrophotometer.

Calculating ATP-Dependent Transport:

ATP-dependent transport = uptake in the presence of ATP – uptake in the presence of AMP.

This product is strictly for laboratory research use only.

