

Neuronostatin-13 (Human, Porcine) - Antibody for Immunohistochemistry

Catalog#	Standard size	Price	Order
H-060-50	100 ul	\$450.00	<input type="button" value="ORDER"/>


Product description

More information

References

Neuronostatin

Image



Human pancreas tissue was stained by Anti-Neuronostatin-13 (H,P) Antibody (Cat.No.: H-060-50)

Protocol for Immunohistochemistry:

Tissue Sample	Human pancreas tissue
Fixative	10% Formalin
Embedding	Paraffin
Negative control	No primary antibody
Pretreatment	No
Blocking	2% Normal Goat Serum
Primary Antibody	Rabbit Anti-Neuronostatin-13 (Human, Porcine) Antibody (Catalog No.: H-060-50)
Optimal Dilution	1:200 (1hour at RT)
Secondary Antibody	Goat anti-Rabbit IgG, Biotinylated (1:400, 30 min)
Amplification	ABC (Vector) (1:400, 30 min)
Detection system	HRP
Substrate	DAB (Sigma), 3 min
Counterstained	Hematoxylin, 30 Sec

Catalog # H-060-50

Family Neuronostatin

Standard Size 100 ul

Species Human, Porcine

Host rabbit

Immunogen human/porcine Neuronostatin-13

Storage Condition For optimal results, use the antibody as soon as possible after reconstitution. Store in lyophilized form unless needed and reconstitute immediately before use. Once reconstituted, the antibody should be stable for a few days at -4°C. For storage up to a few months, prepare small aliquots after reconstitution and freeze at -20°C or -80°C. Repeated freeze thaw cycles should be strictly avoided.