

## DESCRIPTION

<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human Prominin 2. Stains human Prominin 2-transfected cells but not irrelevant transfectants.
<b>Source</b>	Monoclonal Mouse IgG <sub>2B</sub> Clone # 244029
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	NS0 mouse myeloma cell line transfected with human Prominin 2 Ala27-Leu834 Accession # AAM10541
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Flow Cytometry</b>	2.5 µg/10 <sup>6</sup> cells	LNCaP human prostate cancer cell line
<b>Immunocytochemistry</b>	8-25 µg/mL	Immersion fixed PC-3 human prostate cancer cell line

## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

Prominin 2 is a pentaspan membrane glycoprotein predominantly expressed in neuroepithelial cells, hematopoietic stem cells and epithelial cells of the adult kidney and digestive tract (1, 2). Prominin 2 is a 112-kDa glycoprotein structurally related to prominin 1 (CD133). The amino acid identity between prominin 1 and prominin 2 is low (<30%), but they exhibit some redundant functions.

### References:

1. Fargeas, C.A. *et al.* (2003) *J. Biol. Chem.* **278**(10):8586.
2. Fargeas, C.A. *et al.* (2003) *Stem Cells* **21**(4):506.