

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human CDCP1 in direct ELISAs and Western blots. Recognizes an epitope found between aa 33-369 of human CDCP1.
Source	Monoclonal Mouse IgG ₁ Clone # 309137
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human CDCP1 isoform 1 Ala33-Leu666 Accession # NP_073753
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	Recombinant Human CDCP1
Immunoprecipitation	25 µg/mL	Conditioned cell culture medium spiked with Recombinant Human CDCP1, see our available Western blot detection antibodies

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

CDCP1, also known as SIMA135, is a novel 140 kDa type I transmembrane glycoprotein with three CUB protein-protein interaction domains in its 635 aa extracellular region. The 148 aa cytoplasmic region contains canonical phosphorylation sites for Src kinase family members and binding sites for SH3 domains. By alternative splicing, a secreted 310 aa residue form of CDCP1 also exists. The amino-terminal region of approximately 265 aa of the type I membrane CDCP1 can also be proteolytically cleaved. CDCP1 is found on the surface of epithelial and bone marrow-derived stem cells. The extracellular region of human CDCP1 shares 84% aa identity with that of the mouse protein.