MATERIAL DATA SHEET

Suc-Leu-Leu-Val-Tyr-AMC (Suc-LLVY-AMC) Cat # S-280

Fluorogenic substrate for measuring the chymotrypsin-like peptidase activity of the 20S proteasome. The 20S complex is composed of 28 subunits, arranged in an $\alpha_7\beta_7\beta_7\alpha_7$ stoichiometry. Each of the two internal β -type rings harbors three different proteolytically active sites, provided by the amino-terminal residues of three constitutive subunits: β 1 (post-glutamyl peptide hydrolase site), β 2 (trypsin-like site) and β 5 (chymotrypsin-like site).

Product Information

Quantity: 5 mg

Formula: $C_{40}H_{53}N_5O_{10}$ **Formula Weight:** 763.9

Structure:

Physical/Chemical Characteristics

Stock: Soluble at \geq 20 mM in DMSO. For best results, pellet dry compound prior to

reconstitution.

Purity: > 95% by TLC, HPLC. Structure confirmed by NMR.

Use & Storage

Suc-LLVY-AMC is a fluorogenic substrate for measuring the chymotrypsin-

like hydrolyzing activity of the 20S proteasome. Release of AMC

Use: fluorescence can be monitored with an excitation wavelength of 345 nm and

an emission wavelength of 445 nm. Reaction conditions will need to be

optimized for each specific application.

Storage: Store DMSO stock at -20°C. Avoid multiple freeze/thaw cycles.

840 Memorial Drive, Cambridge, MA 02139 Phone: 617-576-2210 FAX: 617-492-3565 www.bostonbiochem.com



Literature

References: Arendt C. S. and Hochstrasser M. (1997) Proc. Natl. Acad. Sci. **94**: 7156

Chen P. and Hochstrasser M. (1996) <u>Cell</u> **86**: 961 Coux O., *et al.* (1996) <u>Ann. Rev. Biochem</u>. **65**: 801 Dick T. P., *et al.* (1998) <u>J. Biol. Chem</u>. **273**: 25637 Stein R.L., *et al.* (1996) <u>Biochem</u>. **5**: 3899

For Laboratory Research Use Only, Not For Use in Humans

Rev: 12/23/2014