# ATR(Ab-428) Antiobdy

Catalog No: #21505

Package Size: #21505-1 50ul #21505-2 100ul #21505-4 25ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

### Description


Product Name	ATR(Ab-428) Antiobdy
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were
	purified by affinity-chromatography using epitope-specific peptide.
Applications	IHC
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of total ATR protein.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around aa.426~430 (G-I-S-P-K) derived from Human ATR.
Target Name	ATR
Other Names	FRP1; MEC1; SCKL; SCKL1;
Accession No.	Swiss-Prot: Q13535NCBI Protein: NP_001175.2
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%
	sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

### **Application Details**

Predicted MW: 250kd

Immunohistochemistry: 1:50~1:100

#### **Images**



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using ATR(Ab-428) Antiobdy #21505(left) or the same antibody preincubated with blocking peptide(right).

## Background

ATR encoded by this gene belongs the PI3/PI4-kinase family, and is most closely related to ATM, a protein kinase encoded by the gene mutated in

ataxia telangiectasia. This protein and ATM share similarity with Schizosaccharomyces pombe rad3, a cell cycle checkpoint gene required for cell cycle arrest and DNA damage repair in response to DNA damage. This kinase has been shown to phosphorylate checkpoint kinase CHK1, checkpoint proteins RAD17, and RAD9, as well as tumor suppressor protein BRCA1. Mutations of this gene are associated with Seckel syndrome. An alternatively spliced transcript variant of this gene has been reported, however, its full length nature is not known. Transcript variants utilizing alternative polyA sites exist.

Zhou, X.Z. et al. (1999) Cell Mol. Life Sci. 56, 788-806.

Pinna, L.A. and Ruzzene, M. (1996) Biochim. Biophys. Acta 1314, 191-225.

Kastan, M.B. and Lim, D.S. (2000) Nat. Rev. Mol. Cell Biol. 1, 179-186.

Shechter, D. et al. (2004) DNA Repair (Amst) 3, 901-908.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.