



ZYMO RESEARCH

THE  
Epigenetics  
COMPANY™

## EZ-96 DNA Methylation-Gold™ Kit (Deep-Well Format)

High-throughput and simplified bisulfite conversion processes

### Highlights

- Complete, high-throughput (96-well) bisulfite conversion of GC-rich DNA in less than 3 hours.
- A coupled heat denaturation/conversion reaction step streamlines the conversion of unmethylated cytosines into uracil.
- DNA precipitations are omitted. Instead, DNA is cleaned and desulphonated in a single step.
- Eluted, ultra-pure DNA is ideal for use in subsequent molecular-based analyses.

Catalog Numbers:  
D5008



Scan with your smart-phone camera to  
view the online protocol/video.



tech@zymoresearch.com



www.zymoresearch.com



Toll Free: (888) 882-9682

# Table of Contents

---

<b>Product Contents</b> .....	<b>01</b>
<b>Introduction to DNA Methylation</b> .....	<b>02</b>
<b>Specifications</b> .....	<b>04</b>
<b>Product Description</b> .....	<b>05</b>
<b>Protocol</b> .....	<b>07</b>
Reagent Preparation .....	<b>07</b>
Sample Processing .....	<b>07</b>
<b>Appendix</b> .....	<b>10</b>
<b>Frequently Asked Questions</b> .....	<b>12</b>
<b>Ordering Information</b> .....	<b>13</b>
<b>Guarantee</b> .....	<b>15</b>

# Product Contents

<b>EZ-96 DNA Methylation-Gold™ Kit</b>	<b>D5008 (2 x 96 Rxns.)</b>	<b>Storage Temperature</b>
CT Conversion Reagent*	2 Bottles	Room Temp.
M-Dilution Buffer	7 ml	Room Temp.
M-Dissolving Buffer	1.2 ml	Room Temp.
M-Binding Buffer	125 ml	Room Temp.
M-Wash Buffer**	2 x 36 ml	Room Temp.
M-Desulphonation Buffer	40 ml	Room Temp.
M-Elution Buffer	8 ml	Room Temp.
Zymo-Spin™ I-96 Binding Plates	2 Plates	Room Temp.
Conversion Plates w/ Pierceable Cover Film	2 Plates/Films	Room Temp.
Collection Plates	2 Plates	Room Temp.
Elution Plates	2 Plates	Room Temp.
Instruction Manual	1	-

\* 9 ml water, 500 µl **M-Dissolving Buffer**, and 3 ml **M-Dilution Buffer** must be added per bottle of **CT Conversion Reagent** prior to use.


\*\* Add 144 ml of 100% ethanol to the 36 ml **M-Wash Buffer** concentrate before use.

# Introduction to DNA Methylation

DNA methylation is a naturally occurring event in both prokaryotic and eukaryotic organisms. In prokaryotes DNA methylation provides a way to protect host DNA from digestion by restriction endonucleases that are designed to eliminate foreign DNA, and in higher eukaryotes DNA methylation functions in the regulation/control of gene expression (1). It has been demonstrated that aberrant DNA methylation is a widespread phenomenon in cancer and may be among the earliest changes to occur during oncogenesis (2). DNA methylation has also been shown to play a central role in gene imprinting, embryonic development, X-chromosome gene silencing, and cell cycle regulation. In many plants and animals, DNA methylation consists of the addition of a methyl group to the fifth carbon position of the cytosine pyrimidine ring via a methyltransferase enzyme (3). The majority of DNA methylation in mammals occurs in 5'-CpG-3' dinucleotides, but other methylation patterns do exist. In fact, about 80 percent of all 5'-CpG-3' dinucleotides in mammalian genomes are found to be methylated, whereas the majority of the twenty percent that remain unmethylated are within promoters or in the first exons of genes.

The ability to detect and quantify DNA methylation efficiently and accurately has become essential for the study of cancer, gene expression, genetic diseases, as well as many other important aspects of biology. To date, a number of methods have been developed to detect/quantify DNA methylation including: high-performance capillary electrophoresis (4) and methylation-sensitive arbitrarily primed PCR (5). However, the most common technique used today remains the bisulfite conversion method (6). This technique involves treating methylated DNA with bisulfite, which converts unmethylated cytosines into uracil. Methylated cytosines remain unchanged during the treatment. Once converted, the methylation profile of the DNA can be determined by PCR amplification followed by DNA sequencing (see next page).

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	
Original DNA with methylated C <sup>m</sup> pG	▶	G	T	T	G	C <sup>m</sup> G	C	T	C	A	C	T	G	C	C	
DNA Sequencing after CT conversion	▶	G	T	T	G	C	G	T	T	T	A	T	T	G	T	T



**DNA sequencing results following bisulfite treatment.** DNA with methylated C<sup>m</sup>pG at nucleotide position #5 was processed using the **EZ DNA Methylation™ Kit**. The recovered DNA was amplified by PCR and then sequenced directly. The methylated cytosine at position #5 remained intact while the unmethylated cytosines at positions #7, 9, 11, 14 and 15 were completely converted into uracil following bisulfite treatment and detected as thymine following PCR.

## References:

1. Costello JF, Plass CJ. *Med. Genet.* 2001; 38(5): 285-303.
2. Stirzaker C. *Cancer Res.* 1997; 57(11): 2229-2237.
3. Adams RL. *Bioessays.* 1995; 17(2): 139-145.
4. Fraga MF, *et al.* *Electrophoresis.* 2000; 21(14): 2990-2994.
5. Gonzalgo ML. *Cancer Res.* 1997; 57(4): 594-599.
6. Frommer M. *Proc. Natl. Acad. Sci. USA.* 1992; 89(5): 1827-1831.


# Specifications

- **DNA Input** – Samples containing 500 pg - 2 µg of DNA. For optimal results, the amount of input DNA should be from 200 to 500 ng.
- **Conversion Efficiency** – > 99% of non-methylated C residues are converted to U; > 99% protection of methylated cytosines.
- **DNA Recovery** – > 75%

# Product Description

The **EZ DNA Methylation-Gold™ Kit** is a refinement of our popular **EZ DNA Methylation™ Kit**. The **EZ DNA Methylation-Gold™ Kit** integrates DNA denaturation and bisulfite conversion processes into one-step. This is accomplished using temperature denaturation to replace chemical denaturation with sodium hydroxide in the previous protocol. Also, the kit has been streamlined for high yield recovery of DNA following DNA bisulfite conversion. Both kits are based on a three-step reaction process between cytosine and sodium bisulfite resulting in cytosine being converted into uracil. The **EZ DNA Methylation-Gold™** and **EZ DNA Methylation™ Kits** share innovative in-column desulphonation technology that eliminates cumbersome DNA precipitation steps while providing researchers consistent results every time. The kits have been designed to minimize template degradation, loss of DNA during treatment and clean-up, and to provide complete conversion of unmethylated cytosines. Recovered DNA is ideal for PCR amplification for downstream analyses including endonuclease digestion, sequencing, microarrays, etc.

An outline comparing the **EZ DNA Methylation-Gold™ Kit** procedure to Zymo Research's other methylation kits is shown below.



Outline of the **EZ DNA Methylation™**, **EZ DNA Methylation-Gold™** and **EZ DNA Methylation-Direct™** Kit procedures.

**Note:** 96-Well spin-plate formats are available for processing larger numbers of samples. Also, MagPrep kits are available (p. 13) for adaptation to liquid handling robots (e.g., Tecan – Freedom EVO®) and automated sample prep.

**Selected EZ DNA Methylation™ Kit Citations:**

1. Ehrich M, *et al.* Nuc. Acids Res. 2007; 35 (5): e29
2. Kaneda M, *et al.* Nature. 2004; 429: 900-903
3. Zhang F, *et al.* Proc. Natl. Acad. Sci. USA. 2007; 104 (11): 4395-4400.
4. Oda M, *et al.* Genes & Dev. 2006; 20: 3382-3394.
5. England RPM, *et al.* Nature Meth. 2005; 2: 1-2.



# Protocol

## Reagent Preparation

### Preparation of CT Conversion Reagent

- ✓ The **CT Conversion Reagent** supplied within this kit is a solid mixture and must be prepared prior to first use. Prepare as follows:
  1. Add 9 ml water, 500  $\mu$ l **M-Dissolving Buffer**, and 3 ml of **M-Dilution Buffer** to a bottle of **CT Conversion Reagent**.
  2. Mix at room temperature with frequent vortexing or shaking for 15 minutes.

**Note:** It is normal to see trace amounts of undissolved reagent in the **CT Conversion Reagent**. Each bottle of **CT Conversion Reagent** is designed for 96 separate DNA treatments.

**Storage:** The **CT Conversion Reagent** is light sensitive, so minimize its exposure to light. For best results, the **CT Conversion Reagent** should be used immediately following preparation. If not used immediately, the **CT Conversion Reagent** solution can be stored overnight at room temperature, one week at 4°C, or up to one month at -20°C. Stored **CT Conversion Reagent** solution must be warmed to 37°C, then vortexed prior to use.

### Preparation of M-Wash Buffer

- ✓ Add 144 ml of 100% ethanol to the 36 ml **M-Wash Buffer** concentrate before use.

## Sample Processing

1. Add 130  $\mu\text{l}$  of the **CT Conversion Reagent** to 20  $\mu\text{l}$ \* of each DNA sample in a **Conversion Plate**. If the volume of the DNA sample is less than 20  $\mu\text{l}$ , make up the difference with water. Mix the samples by pipetting up and down.
2. Seal the plate with the provided film. Transfer the **Conversion Plate** to a thermal cycler and perform the following steps:
  1. 98°C for 10 minutes
  2. 64°C for 2.5 hours
  3. 4°C storage for up to 20 hours

**Note:** The 4°C storage step is *optional*. For some samples, alternative parameters may yield improved results (see Appendix). If you have been using this kit with good results using different reaction conditions than described above, you can continue using those same conditions.

3. Add 600  $\mu\text{l}$  of **M-Binding Buffer** to the wells of a **Zymo-Spin™ I-96 Binding Plate** mounted on a **Collection Plate**.

**Note:** The capacity of each well of the Binding Plate is 1.1 ml. The capacity of each well of the Collection Plate is 800  $\mu\text{l}$ . Empty the Collection Plate whenever necessary to prevent contamination of the Binding Plate contents by the flow-through.

4. Transfer the samples from the **Conversion Plate** (Step 2) to the wells of the **ZymoSpin™ I-96 Binding Plate**. Mix by pipetting up and down.
5. Centrifuge at  $\geq 3,000 \times g$  (5,000  $\times g$  max.) for 5 minutes. Discard the flow-through.
6. Add 400  $\mu\text{l}$  of **M-Wash Buffer** to each well of the plate. Centrifuge at  $\geq 3,000 \times g$  for 5 minutes.
7. Add 200  $\mu\text{l}$  of **M-Desulphonation Buffer** to each well and allow the plate to stand at room temperature (20-30°C) for 15-20 minutes. After the incubation, centrifuge at  $\geq 3,000 \times g$  for 5 minutes. Discard the flow-through.

---

\*For DNA volumes  $>20 \mu\text{l}$ , an adjustment needs to be made during the preparation of the **CT Conversion Reagent**. The amount of water is decreased 1 ml for each 10  $\mu\text{l}$  increase in DNA sample volume. For example, for 40  $\mu\text{l}$  DNA samples, 7 ml of water is added to make the **CT Conversion Reagent**. The volume of **CT Conversion Reagent** added to the sample must also be decreased by the same volume as the sample is increased, total reaction volume remains 150  $\mu\text{l}$ . The maximum DNA sample volume to be used for each conversion reaction is 45  $\mu\text{l}$ . Do not adjust the volumes of either the **M-Dissolving Buffer** or **M-Dilution Buffer**.

8. Add 400  $\mu\text{l}$  of **M-Wash Buffer** to each well of the plate. Centrifuge at  $\geq 3,000 \times g$  for 5 minutes. Discard the flow-through. Add another 400  $\mu\text{l}$  of **M-Wash Buffer** and centrifuge for 10 minutes.
9. Place the **Zymo-Spin™ I-96 Binding Plate** onto an **Elution Plate**. Add 15  $\mu\text{l}$  of **M-Elution Buffer** directly to each well. After 5 minutes, centrifuge at  $\geq 3,000 \times g$  for 3 minutes to elute the DNA.

**Note:** Alternatively, water or TE (pH  $\geq 6.0$ ) can be used for elution if required for your experiments.

The DNA is ready for immediate analysis or can be stored at or below  $-20^{\circ}\text{C}$  for later use. For long term storage, store at or below  $-70^{\circ}\text{C}$ . We recommend using 1-4  $\mu\text{l}$  of eluted DNA for each PCR, however, up to 15  $\mu\text{l}$  can be used if necessary. The elution volume can be  $> 15 \mu\text{l}$  depending on the requirements of your experiments, but small elution volumes will yield more concentrated DNA.

# Appendix

## Bisulfite Conversion and PCR Optimization

1. **Reaction Conditions:** The reaction conditions given in Step 2 of the Protocol will generate consistent results for both easy and difficult to convert template DNAs including those that are GC rich. However, the two protocols provided below (alternative 1 & 2) may yield better results in PCR amplification of longer DNA fragments. However, should the DNA template have >80% GC composition, then these conditions may result in incomplete template cytosine to uracil conversion.

### Alternative 1:

1. 98°C for 10 minutes
  2. 53°C for 30 minutes
  3. 53°C for 6 minutes
  4. 37°C for 30 minutes
  5. 4°C storage
- } 8 cycles

### Alternative 2:

1. 98°C for 10 minutes
2. 53°C for 4 hours
3. 4°C storage

2. **Bisulfite Conversion of Double Stranded DNA Templates.** The following illustrates what occurs to a DNA template during bisulfite conversion.

Template:                    **A:** 5' -GACCGTTCCAGGTCCAGCAGTGCGCT-3'  
                                     **B:** 3' -CTGGCAAGGTTCCAGGTCGTCACGCGA-5'

Bisulfite Converted:    **A:** 5' -GATCGTTTTAGGTTTAGTAGTGCGTT-3'  
                                     **B:** 3' -TTGGCAAGGTTTAGGTTTATGCGA-5'

**Note:** Methylated "C" is underlined in the examples.

**Note:** Following bisulfite conversion, the strands are no longer complementary.

- 3. PCR Primer Design.** Generally, primers 26 to 32 bases are required for amplification of bisulfite converted DNA. In general, all Cs should be treated as Ts for primer design purposes, unless they are in a CpG context. See example below.

Bisulfite Converted: **A:** 5' -GATCGTTTTAGGTTTAGTAGTGCGTT-3'  
Primers: Reverse: 3' -ATCATCACRCAA-5' R= G/A  
: Forward: 5' -GATYGTTTTAGGT-3' Y= C/T

**Note:** Only one strand (**A**) is amplified by a given primer set. Only the reverse primer binds to the converted DNA, the forward primer will bind the strand generated by the reverse primer. If the primer contains CpG dinucleotides with uncertain methylation status, then mixed bases with C and T (or G and A) can be used. Usually, there should be no more than one mixed position per primer and it should be located toward the 5' end of the primer. It is not recommended to have mixed bases located at the 3' end of the primer.

Zymo Research provides primer design assistance with its Bisulfite Primer Seeker Program, available at:

[www.zymoresearch.com/tools/bisulfite-primer-seeker](http://www.zymoresearch.com/tools/bisulfite-primer-seeker)

- 4. Amount of DNA Required for Bisulfite Conversion.** The minimal amount of human or mouse genomic DNA required for bisulfite treatment and subsequent PCR amplification is 100 pg. The optimal amount of DNA per bisulfite treatment is 200 to 500 ng. Although, up to 2 µg of DNA can be processed, it should be noted that high input levels of DNA may result in incomplete bisulfite conversion for some GC-rich regions.
- 5. PCR Conditions.** Usually, 35 to 40 cycles are required for successful PCR amplification of bisulfite converted DNA. Optimal amplicon size should be between 150-300 bp; however larger amplicons (up to 1 kb) can be generated by optimizing the PCR conditions. Annealing temperatures between 55-60°C typically work well.

As most non-methylated cytosine residues are converted into uracil, the bisulfite-treated DNA usually is AT-rich and has low GC composition. Non-specific PCR amplification is relatively common with bisulfite treated DNA due to its AT-rich nature. PCR using “hot start” polymerases is strongly recommended for the amplification of bisulfite-treated DNA.

**Note:** **ZymoTag™** is a “hot start” DNA polymerase specifically designed for the amplification of bisulfite treated DNA.

## Frequently Asked Questions

**Q: Should the input DNA be dissolved in TE, water, or some other buffer prior to its conversion?**



*A: Water, TE or modified TE buffers can be used to dissolve the DNA and do not interfere with the conversion process.*

**Q: Which Taq polymerase(s) do you recommend for PCR amplification of converted DNA?**

*A: We recommend a “hot start” DNA polymerase (e.g., ZymoTaq™ DNA Polymerase).*

**Q: Why are there two different catalog numbers for the EZ-96 DNA Methylation™ Kit?**

*A: The two different catalog numbers are used to differentiate between the binding plates that are included in the kit. Deep and shallow-well binding plates are available to accommodate most rotors and microplate carriers. Below is a comparison of the two binding plates.*

		
<b>Binding Plate</b>	<b>Silicon-A™ Plate</b>	<b>Zymo-Spin™ I-96 Plate</b>
<b>Style</b>	Shallow-Well	Deep-Well
<b>Height of Binding Plate</b>	19 mm (0.75 inches)	35 mm (1.38 inches)
<b>Binding Plate/Collection Plate Assembly</b>	43 mm (1.69 inches)	60 mm (2.36 inches)
<b>Binding Cap./Minimum Elution Volume</b>	5 µg/30 µl	5 µg/15 µl
<b>Catalog Numbers</b>	<b>D5007</b>	<b>D5008</b>

# Ordering Information

Product Description	Catalog No.	Size
EZ DNA Methylation-Gold™ Kit	D5005	50 Rxns.
	D5006	200 Rxns.
EZ-96 DNA Methylation-Gold™ Kit (Shallow-Well)	D5007	2 x 96 Rxns.
EZ-96 DNA Methylation-Gold™ Kit (Deep-Well)	D5008	2 x 96 Rxns.
EZ-96 DNA Methylation-Gold™ MagPrep*	D5046	4 x 96 Rxns.
	D5047	8 x 96 Rxns.

\* MagPrep kits are adaptable to liquid handling robots (e.g., Tecan – Freedom EVO®) making them ideal for automated sample prep.

Individual Kit Components	Catalog No.	Amount
CT Conversion Reagent	D5001-1	1 Tube
	D5003-1	1 Bottle
M-Dilution Buffer	D5005-2	1.5 ml
	D5006-2	7 ml
M-Binding Buffer	D5005-3	30 ml
	D5006-3	125 ml
	D5040-3	250 ml
M-Wash Buffer	D5001-4	6 ml
	D5002-4	24 ml
	D5007-4	36 ml
	D5040-4	72 ml
M-Desulphonation Buffer	D5001-5	10 ml
	D5002-5	40 ml
	D5040-5	80 ml
M-Elution Buffer	D5001-6	1 ml
	D5002-6	4 ml
	D5007-6	8 ml
	D5041-6	40 ml
M-Dissolving Buffer	D5005-6	500 µl
	D5006-6	1.2 ml
Zymo-Spin™ IC Columns (capped)	C1004-50	50 Pack
	C1004-250	250 Pack

<b>Collection Tubes</b>	C1001-50 C1001-500 C1001-1000	50 Pack 500 Pack 1,000 Pack
<b>MagBinding Beads</b>	D4100-5-3 D4100-5-8 D4100-5-16	6 ml 8 ml 16 ml
<b>Zymo-Spin™ I-96 Binding Plates</b>	C2004	2 Plates
<b>Silicon-A™ Binding Plates</b>	C2001	2 Plates
<b>Conversion Plates w/ Pierceable Cover Film</b>	C2005	2 Plates /Films
<b>Collection Plates</b>	C2002	2 Plates
<b>Elution Plates</b>	C2003	2 Plates





**100% satisfaction guarantee on all Zymo Research products,  
or your money back.**

Zymo Research is committed to simplifying your research with quality products and services. If you are dissatisfied with this product for any reason, please call 1(888) 882-9682.

---

Integrity of kit components is guaranteed for up to one year from date of purchase. Reagents are routinely tested on a lot-to-lot basis to ensure they provide the highest performance and reliability.

This product is for research use only and should only be used by trained professionals. It is not for use in diagnostic procedures. Some reagents included with this kit are irritants. Wear protective gloves and eye protection. Follow the safety guidelines and rules enacted by your research institution or facility.

EZ DNA Methylation-Gold™ Kit technologies are patent pending.

Use of Methylation Specific PCR (MSP) is protected by US Patents 5,786,146 & 6,017,704 & 6,200,756 & 6,265,171 and International Patent WO 97/46705. No license under these patents to use the MSP process is conveyed expressly or by implication to the purchaser by the purchase of this product.

™ Trademarks of Zymo Research Corporation



ZYMO RESEARCH

The **BEAUTY** of **SCIENCE** is to Make Things **SIMPLE**®



[tech@zymoresearch.com](mailto:tech@zymoresearch.com)



[www.zymoresearch.com](http://www.zymoresearch.com)



Toll Free: (888) 882-9682