
Gliadin Peptide (4F3) Antibody

Subcategory: Mouse Monoclonal Antibody

Cat. No.: 250438

Unit: 0.2 mg

Description:

Celiac disease is associated with a CD4+ T-cell response to epitopes of gliadin presented by HLA-DQ2 or -DQ8 class II MHC molecules. These epitopes are present in a 33-mer peptide of wheat alpha-gliadin, residues 56-88, which is resistant to digestion and forms a substrate for tissue transglutaminase (TG2), generating the glutamic acid residues essential for binding to HLA-DQ2. The immunogen corresponds to a deamidated form of a region that includes the T-cell epitopes, including the immunodominant PQPQLPY region and two PXPQP motifs associated with binding to IgA from patients with celiac disease. Complete homology exists between residues 63-73, 70-80 and 77-87 of wheat alpha gliadin. This antibody cross-reacts fully with the non-deamidated peptide KLQPFQPQLPYQPQ

Isotype: Mouse IgG2a/k

Applications: E

Species Reactivity: PI

Format: Each vial contains 0.2 mg IgG in 0.2 ml (1 mg/ml) of PBS pH7.4 with 0.09% sodium azide. Antibody was purified by Protein-A/G affinity chromatography.

Alternate Names: Alpha gliadin

Accession No.: Q41531

Antigen: Gliadin-related peptide Lys57-Glu65-[alpha-gliadin (58-73)] (KLQPFQPPELPYPQPQ) coupled to immunogenic carrier protein.

Application Notes: Product Cat. No. 250438 reacts equally strong in ELISA with the deamidated gliadin-related peptide (KLQPFQPPELPYPQPQ) and the non-deamidated gliadin-related peptide (KLQPFQPQLPYQPQ) coated directly in microplate. E: 1:20,000

Storage: Store at 4°C. Product is guaranteed 6 months from the date of shipment.

For research use only, not for diagnostic or therapeutic procedures.