

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

## **Datasheet**

## FGR monoclonal antibody (M07), clone 4C12

Catalog Number: H00002268-M07

Regulatory Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody

raised against a partial recombinant FGR.

Clone Name: 4C12

**Immunogen:** FGR (AAH64382, 1 a.a. ~ 90 a.a) partial recombinant protein with GST tag. MW of the GST tag

alone is 26 KDa.

## Sequence:

MGCVFCKKLEPVATAKEDAGLEGDFRSYGAADHYGP DPTKARPASSFAHIPNYSNFSSQAINPGFLDSGTIRGV SGIGVTLFIALYDYEA

Host: Mouse

Reactivity: Human

Applications: ELISA, IHC-P, S-ELISA, WB-Re

(See our web site product page for detailed applications

information)

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Isotype: IgG2a Kappa

Storage Buffer: In 1x PBS, pH 7.4

Storage Instruction: Store at -20°C or lower. Aliquot to

avoid repeated freezing and thawing.

Entrez GenelD: 2268

Gene Symbol: FGR

Gene Alias: FLJ43153, MGC75096, SRC2, c-fgr,

c-src2, p55c-fgr, p58c-fgr

**Gene Summary:** This gene is a member of the Src family of protein tyrosine kinases (PTKs). The encoded protein contains N-terminal sites for myristylation and

palmitylation, a PTK domain, and SH2 and SH3 domains which are involved in mediating protein-protein interactions with phosphotyrosine-containing and proline-rich motifs, respectively. The protein localizes to plasma membrane ruffles, and functions as a negative regulator of cell migration and adhesion triggered by the beta-2 integrin signal transduction pathway. Infection with Epstein-Barr virus results in the overexpression of this gene. Multiple alternatively spliced variants, encoding the same protein, have been identified. [provided by RefSeq]