

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

Datasheet

PSMD5 monoclonal antibody (M01), clone 3E2

Catalog Number: H00005711-M01

Regulatory Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody raised against a partial recombinant PSMD5.

Clone Name: 3E2

 $\label{eq:mmunogen: PSMD5 (AAH14478, 405 a.a. $\sim 504 a.a)} partial recombinant protein with GST tag. MW of the$

GST tag alone is 26 KDa.

Sequence:

QPFPELHCAALKVFTAIANQPWAQKLMFNSPGFVEYV VDRSVEHDKASKDAKYELVKALANSKTIAEIFGNPNYL RLRTYLSEGPYYVKPVSTTAVEGAE

Host: Mouse

Reactivity: Human

Applications: ELISA, IHC-P, S-ELISA, WB-Ce, WB-Re (See our web site product page for detailed applications

information)

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Isotype: IgG1 Kappa

Storage Buffer: In 1x PBS, pH 7.4

Storage Instruction: Store at -20°C or lower. Aliquot to

avoid repeated freezing and thawing.

Entrez GenelD: 5711

Gene Symbol: PSMD5

Gene Alias: KIAA0072, MGC23145, S5B

Gene Summary: The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28

non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a non-ATPase subunit of the 19S regulator base. [provided by RefSeq]