

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

Datasheet

PTPRC monoclonal antibody, clone MEM-55

Catalog Number: MAB0888

Regulatory Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody

raised against native PTPRC.

Clone Name: MEM-55

Immunogen: Native purified PTPRC from human

thymocytes and T lymphocytes.

Host: Mouse

Theoretical MW (kDa): 180-240

Reactivity: Human, Primates

Applications: Flow Cyt, IHC-P, IP, WB

(See our web site product page for detailed applications

information)

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Specificity: This antibody recognizes a siliadase-sensitive epitope of CD45RB, a 180-240 KDa single chain type I membrane glycoprotein, variant of CD45 (CD45RB isoform). CD45RB is expressed on a subset of T lymphocytes, B lymphocytes, monocytes, macrophages, granulocytes and dendritic cells.

Form: Liquid

Isotype: IgG1

Recommend Usage: The optimal working dilution

should be determined by the end user.

Storage Buffer: In PBS, pH 7.4 (0.09% sodium azide)

Storage Instruction: Store at 4°C. Do not freeze. Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 5788

Gene Symbol: PTPRC

Gene Alias: B220, CD45, CD45R, GP180, LCA, LY5,

T200

Gene Summary: The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP contains an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains, and thus belongs to receptor type PTP. This gene is specifically expressed in hematopoietic cells. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen receptor complexes, or by activating various Src family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases, and thus functions as a regulator of cytokine receptor signaling. Four alternatively spliced transcripts variants of this gene, which encode distinct isoforms, have been reported. [provided by RefSeq]