

9F, No. 108, Jhouzih St., Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

## **Datasheet**

## SLC5A5 monoclonal antibody, clone FP5 (FITC)

Catalog Number: MAB11506

Regulatory Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody

raised against recombinant of SLC5A5.

Clone Name: FP5

**Immunogen:** Recombinant mannose fusion protien corresponding to amino acids 468-643 of human

SLC5A5.

Host: Mouse

Reactivity: Human, Mouse, Rat

Applications: IF, IHC, WB

(See our web site product page for detailed applications

information)

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

**Specificity:** Mapped to amino acids 625-643 of hNIS. Apparent mol. wt of 97 kD, non-glycosylated version at 68 kD. Other minor bands associated with hNIS at 160 kDa, and degradation products at ~30 kDa, and ~15kDa.

Form: Liquid

Conjugation: FITC

Purification: Protein G purification

Isotype: IgG1, kappa

Recommend Usage: Immunofluorescence

Immunohistochemistry Western blot (1:1000)

The optimal working dilution should be determined by

the end user.

Storage Buffer: In PBS, pH 7.4 (50% glycerol)

Storage Instruction: Store at 4°C.

Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 6528

Gene Symbol: SLC5A5

Gene Alias: NIS

Gene Summary: This gene encodes a member of the sodium glucose cotransporter family. The encoded protein is responsible for the uptake of iodine in tissues such as the thyroid and lactating breast tissue. The iodine taken up by the thyroid is incorporated into the metabolic regulators triiodothyronine (T3) and tetraiodothyronine (T4). Mutations in this gene are

associated with thyroid dyshormonogenesis 1]