

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

## Datasheet

## ITGAL monoclonal antibody, clone HI111 (PerCP)

Catalog Number: MAB13793

Regulatory Status: For research use only (RUO)

**Product Description:** Mouse monoclonal antibody raised against native human ITGAL.

Clone Name: HI111

Immunogen: Human CD11a.

Host: Mouse

Theoretical MW (kDa): 180

Reactivity: Human

**Applications:** Blocking, Flow Cyt, IHC-Fr, WB (See our web site product page for detailed applications information)

**Protocols:** See our web site at http://www.abnova.com/support/protocols.asp or product page for detailed protocols

Form: Liquid

Conjugation: PerCP

Purification: Protein A/G purification

**Purity:** >90%

Isotype: IgG1, kappa

Recommend Usage: Blocking Flow Cytometry (20 uL/10<sup>6</sup> cells) Immunohistochemistry (Frozen sections) Western Blot The optimal working dilution should be determined by the end user.

**Storage Buffer:** In PBS, pH 7.4 (protein stabilizer, 0.09% sodium azide).

**Storage Instruction:** Store in the dark at 4°C. Avoid prolonged exposure to light.

Entrez GenelD: 3683

Gene Symbol: ITGAL

Gene Alias: CD11A, LFA-1, LFA1A

**Gene Summary:** ITGAL encodes the integrin alpha L chain. Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain. This I-domain containing alpha integrin combines with the beta 2 chain (ITGB2) to form the integrin lymphocyte function-associated antigen-1 (LFA-1), which is expressed on all leukocytes. LFA-1 plays a central role in leukocyte intercellular adhesion through interactions with its ligands, ICAMs 1-3 (intercellular adhesion molecules 1 through 3), and also functions in lymphocyte costimulatory signaling. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]