

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

## Datasheet

## CR1 monoclonal antibody, clone UJ11 (FITC)

Catalog Number: MAB13883

Regulatory Status: For research use only (RUO)

**Product Description:** Mouse monoclonal antibody raised against human CR1.

Clone Name: UJ11

Immunogen: Human cells of the monocyte lineage.

Host: Mouse

Theoretical MW (kDa): 280

Reactivity: Human

**Applications:** Flow Cyt (See our web site product page for detailed applications information)

**Protocols:** See our web site at http://www.abnova.com/support/protocols.asp or product page for detailed protocols

Form: Liquid

Conjugation: FITC

Purification: Protein A/G purification

Purity: >90%

Isotype: IgG1

**Recommend Usage:** Flow Cytometry (20 uL/10<sup>6</sup> cells) The optimal working dilution should be determined by the end user.

**Storage Buffer:** In PBS, pH 7.4 (protein stabilizer, 0.09% sodium azide).

**Storage Instruction:** Store in the dark at 4°C. Avoid prolonged exposure to light.

Entrez GenelD: 1378

Gene Symbol: CR1

Gene Alias: C3BR, CD35, KN

Gene Summary: This gene is a member of the receptors of complement activation (RCA) family and is located in the 'cluster RCA' region of chromosome 1. The gene encodes a monomeric single-pass type I found membrane glycoprotein on erythrocytes, leukocytes, glomerular podocytes, and splenic follicular dendritic cells. The Knops blood group system is a system of antigens located on this protein. The protein mediates cellular binding to particles and immune complexes that have activated complement. Decreases in expression of this protein and/or mutations in its gene have been associated with gallbladder carcinomas, mesangiocapillary glomerulonephritis, systemic lupus erythematosus and sarcoidosis. Mutations in this gene have also been associated with a reduction in Plasmodium falciparum rosetting, conferring protection against severe malaria. Alternate allele-specific splice variants, encoding different isoforms, have been characterized. Additional allele specific isoforms, including a secreted form, have been described but have not been fully characterized. [provided by RefSeq]