

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

Datasheet

CD44 monoclonal antibody, clone HI44a (FITC)

Catalog Number: MAB13922

Regulatory Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody

raised against human CD44.

Clone Name: HI44a

Immunogen: Human leukocytes of a patient suffering

from LGL type leukemia.

Host: Mouse

Theoretical MW (kDa): 80-95

Reactivity: Human

Applications: Flow Cyt

(See our web site product page for detailed applications

information)

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Form: Liquid

Conjugation: FITC

Purification: Protein A/G purification

Purity: >90%

Isotype: IgG2a

Recommend Usage: Flow Cytometry (20 uL/10⁶ cells) The optimal working dilution should be determined by

the end user.

Storage Buffer: In PBS, pH 7.4 (protein stabilizer,

0.09% sodium azide).

Storage Instruction: Store in the dark at 4°C. Avoid

prolonged exposure to light.

Entrez GenelD: 960

Gene Symbol: CD44

Gene Alias: CDW44, CSPG8, ECMR-III, HCELL, IN, LHR, MC56, MDU2, MDU3, MGC10468, MIC4,

MUTCH-I, Pgp1

Gene Summary: The protein encoded by this gene is a cell-surface glycoprotein involved in cell-cell interactions, cell adhesion and migration. It is a receptor for hyaluronic acid (HA) and can also interact with other ligands, such as osteopontin, collagens, and matrix metalloproteinases (MMPs). This protein participates in a wide variety of cellular functions including lymphocyte activation, recirculation and homing, hematopoiesis, and tumor metastasis. Transcripts for this gene undergo complex alternative splicing that results in many functionally distinct isoforms, however, the full length nature of some of these variants has not been determined. Alternative splicing is the basis for the structural and functional diversity of this protein, and may be related to tumor metastasis. [provided by RefSeq]