

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

Datasheet

Mme monoclonal antibody, clone LT10 (FITC)

Catalog Number: MAB4348

Regulatory Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody

raised against native Mme.

Clone Name: LT10

Immunogen: Mative purified Mme from mouse NALM-6

leukemia pre-B cell line.

Host: Mouse

Theoretical MW (kDa): 100

Reactivity: Human

Applications: Flow Cyt

(See our web site product page for detailed applications

information)

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Specificity: This antibody reacts with CD10 antigen (CALLA - Common acute lymphatic leukemia antigen), a

100 KDa type II integral membrane protein.

Form: Liquid

Conjugation: FITC

Isotype: IgG1

Recommend Usage: Flow Cytometry (20 ul in human

blood cells 100 ul in whole blood or 10⁶ cells in a

suspension)

The optimal working dilution should be determined by

the end user.

Storage Buffer: In PBS (0.2% BSA, 0.09% sodium

azide)

Storage Instruction: Store in the dark at 4°C. Do not

freeze.

Avoid prolonged exposure to light.

Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 17380

Gene Symbol: Mme

Gene Alias: 6030454K05Rik, C85356, CALLA, CD10,

NEP

References:

- 1. Differential expression of CD10 in prostate cancer and its clinical implication. Dall'Era MA, True LD, Siegel AF, Porter MP, Sherertz TM, Liu AY. BMC Urol. 2007 Mar 2:7:3.
- 2. CD10 expression by fusiform stromal cells in nasopharyngeal carcinoma correlates with tumor progression. Braham H, Trimeche M, Ziadi S, Mestiri S, Mokni M, Amara K, Hachana M, Sriha B, Korbi S. Virchows Arch. 2006 Aug;449(2):220-4. Epub 2006 May 4.
- 3. Expression of CD10 in basal cell carcinoma. Yada K, Kashima K, Daa T, Kitano S, Fujiwara S, Yokoyama S. Am J Dermatopathol. 2004 Dec;26(6):463-71.