

Datasheet

CD27 monoclonal antibody, clone LT27 (PE)

Catalog Number: MAB4400

Regulatory Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody raised against native CD27.

Clone Name: LT27

Immunogen: Native purified CD27 from human peripheral blood lymphocytes.

Host: Mouse

Theoretical MW (kDa): 50-55

Reactivity: Human

Applications: Flow Cyt

(See our web site product page for detailed applications information)

Protocols: See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Specificity: This antibody reacts with CD27 (T14), a 50-55 kDa type I transmembrane glycoprotein (member of the TNF-receptor superfamily) expressed on medullary thymocytes, peripheral T lymphocytes, some B lymphocytes and NK cells.

Form: Liquid

Conjugation: PE

Isotype: IgG2a

Recommend Usage: Flow Cytometry (20 ul in human blood cells 100 ul in whole blood or 10⁶ cells in a suspension)

The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS (0.2% BSA, 0.09% sodium azide)

Storage Instruction: Store in the dark at 4 °C. Do not freeze.

Avoid prolonged exposure to light.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 939

Gene Symbol: CD27

Gene Alias: MGC20393, S152, T14, TNFRSF7, Tp55

Gene Summary: The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is required for generation and long-term maintenance of T cell immunity. It binds to ligand CD70, and plays a key role in regulating B-cell activation and immunoglobulin synthesis. This receptor transduces signals that lead to the activation of NF-kappaB and MAPK8/JNK. Adaptor proteins TRAF2 and TRAF5 have been shown to mediate the signaling process of this receptor. CD27-binding protein (SIVA), a proapoptotic protein, can bind to this receptor and is thought to play an important role in the apoptosis induced by this receptor. [provided by RefSeq]

References:

1. Blockade of protease-activated receptors on T cells correlates with altered proteolysis of CD27 by gingipains of *Porphyromonas gingivalis*. Yun LW, Decarlo AA, Hunter N. Clin Exp Immunol. 2007 Nov;150(2):217-29.
2. Expression of soluble CD27 and interleukins-8 and -10 in B-cell chronic lymphocytic leukemia: correlation with disease stage and prognosis. Kara IO, Sahin B, Gunesacar R. Adv Ther. 2007 Jan-Feb;24(1):29-40.
3. CD27 in B-cell chronic lymphocytic leukemia. Cellular expression, serum release and correlation with other soluble molecules belonging to nerve growth factor receptors (NGFr) superfamily. Molica S, Vitelli G, Levato D, Crispino G, Dell'Olio M, Dattilo A, Matera R, Gandolfo GM, Musto P. Haematologica. 1998 May;83(5):398-402.