

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

## **Datasheet**

## HLA-G monoclonal antibody, clone 2A12 (FITC)

Catalog Number: MAB4488

Regulatory Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody

raised against synthetic peptide HLA-G.

Clone Name: 2A12

**Immunogen:** A synthetic peptide (conjugated with OVA) corresponding to C-terminus of human soluble HLA-G5

and HLA-G6.

Host: Mouse

Reactivity: Human

Applications: Flow Cyt, IHC-Fr, IHC-P, WB

(See our web site product page for detailed applications

information)

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

**Specificity:** This antibody allows to discriminate between soluble HLA-G protein yielded by shedding from membrane-bound HLA-G forms that do not contain the intron 4-encoded epitope and soluble HLA-G5/HLA-G6 produced from intron 4-retaining alternatively spliced mRNAs. HLA-G belongs to the MHC Class I molecules (MHC Class Ib; nonclassical) and it is expressed on the surface of trophoblast cells.

Form: Liquid

Conjugation: FITC

Concentration: 1 mg/mL

Isotype: IgG1

**Recommend Usage:** Flow Cytometry (5 ug/mL) The optimal working dilution should be determined by

the end user.

Storage Buffer: In PBS, pH 7.4 (0.09% sodium azide)

Storage Instruction: Store in the dark at 4°C. Do not

freeze.

Avoid prolonged exposure to light.

Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 3135

Gene Symbol: HLA-G

Gene Alias: MHC-G

Gene Summary: HLA-G belongs to the HLA class I heavy chain paralogues. This class I molecule is a heterodimer consisting of a heavy chain and a light chain (beta-2 microglobulin). The heavy chain is anchored in the membrane. HLA-G is expressed on fetal derived placental cells. The heavy chain is approximately 45 kDa and its gene contains 8 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the alpha1 and alpha2 domain, which both bind the peptide, exon 4 encodes the alpha3 domain, exon 5 encodes the transmembrane region, and exon 6 encodes the cytoplasmic tail. [provided by RefSeq]