

## Datasheet

### HLA-G monoclonal antibody, clone MEM-G/11 (FITC)

**Catalog Number:** MAB4520

**Regulatory Status:** For research use only (RUO)

**Product Description:** Mouse monoclonal antibody raised against recombinant HLA-G.

**Clone Name:** MEM-G/11

**Immunogen:** Recombinant protein corresponding to human HLA-G.

**Host:** Mouse

**Reactivity:** Human

**Applications:** ELISA, Flow Cyt, IP  
(See our web site product page for detailed applications information)

**Protocols:** See our web site at  
<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Specificity:** This antibody recognizes HLA-G1 antigen. HLA-G belongs to the MHC Class I molecules (MHC Class Ib; nonclassical) and it is expressed on the surface of trophoblast cells.

**Form:** Liquid

**Conjugation:** FITC

**Concentration:** 1 mg/mL

**Isotype:** IgG1

**Recommend Usage:** Flow Cytometry (1:500)  
The optimal working dilution should be determined by the end user.

**Storage Buffer:** In PBS, pH 7.4 (0.09% sodium azide)

**Storage Instruction:** Store in the dark at 4°C. Do not freeze.

Avoid prolonged exposure to light.  
Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 3135

**Gene Symbol:** HLA-G

**Gene Alias:** MHC-G

**Gene Summary:** HLA-G belongs to the HLA class I heavy chain paralogues. This class I molecule is a heterodimer consisting of a heavy chain and a light chain (beta-2 microglobulin). The heavy chain is anchored in the membrane. HLA-G is expressed on fetal derived placental cells. The heavy chain is approximately 45 kDa and its gene contains 8 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the alpha1 and alpha2 domain, which both bind the peptide, exon 4 encodes the alpha3 domain, exon 5 encodes the transmembrane region, and exon 6 encodes the cytoplasmic tail. [provided by RefSeq]

#### References:

1. Characterization of monoclonal antibodies recognizing HLA-G or HLA-E: new tools to analyze the expression of nonclassical HLA class I molecules. Menier C, Saez B, Horejsi V, Martinozzi S, Krawice-Radanne I, Bruel S, Le Danff C, Reboul M, Hilgert I, Rabreau M, Larrad ML, Pla M, Carosella ED, Rouas-Freiss N. Hum Immunol. 2003 Mar;64(3):315-26.
2. Disulfide bond-mediated dimerization of HLA-G on the cell surface. Boyson JE, Erskine R, Whitman MC, Chiu M, Lau JM, Koopman LA, Valter MM, Angelisova P, Horejsi V, Strominger JL. Proc Natl Acad Sci U S A. 2002 Dec 10;99(25):16180-5. Epub 2002 Nov 26.