

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

Datasheet

CD8 monoclonal antibody, clone MEM-31 (PerCP)

Catalog Number: MAB4602

Regulatory Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody

raised against native CD8.

Clone Name: MEM-31

Immunogen: Native purified CD8 from Crude thymus

membrane fraction.

Host: Mouse

Reactivity: Human

Applications: Flow Cyt

(See our web site product page for detailed applications

information)

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Specificity: This antibody recognizes a conformationally-dependent epitope of CD8, a cell surface glycoprotein found on most cytotoxic T lymphocytes that mediates efficient cell-cell interactions within the immune system. CD8 is a disulfide-linked dimer and exists as a CD8 alpha/alpha homodimer or CD8 alpha/beta heterodimer (each monomer approx. 32-34 KDa).

This antibody does not react with formaldehyde-fixed cells; negative in Western Blot application.

Form: Liquid

Conjugation: PerCP

Isotype: IgG2a

Recommend Usage: Flow Cytometry (10 ul in human blood cells 100 ul in whole blood or 10⁶ cells in a suspension)

The optimal working dilution should be determined by

the end user.

Storage Buffer: In PBS (0.2% BSA, 0.09% sodium

azide)

Storage Instruction: Store in the dark at 4°C. Do not

freeze.

Avoid prolonged exposure to light.

Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 925|926

Gene Symbol: CD8A

Gene Alias: CD8, Leu2, MAL, p32

References:

- 1. Coreceptor CD8-driven modulation of T cell antigen receptor specificity. van den Berg HA, Wooldridge L, Laugel B, Sewell AK. J Theor Biol. 2007 Nov 21;249(2):395-408. Epub 2007 Aug 8.
- 2. CD8 Raft localization is induced by its assembly into CD8alpha beta heterodimers, Not CD8alpha alpha homodimers. Pang DJ, Hayday AC, Bijlmakers MJ. J Biol Chem. 2007 May 4;282(18):13884-94. Epub 2007 Mar 6.
- 3. Monoclonal antibodies against human leucocyte antigens. II. Antibodies against CD45 (T200), CD3 (T3), CD43, CD10 (CALLA), transferrin receptor (T9), a novel broadly expressed 18-kDa antigen (MEM-43) and a novel antigen of restricted expression (MEM-74). Horejsi V, Angelisova P, Bazil V, Kristofova H, Stoyanov S, Stefanova I, Hausner P, Vosecky M, Hilgert I. Folia Biol (Praha). 1988;34(1):23-34.