

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

## **Datasheet**

## TFRC monoclonal antibody, clone MEM-75 (FITC)

Catalog Number: MAB5015

Regulatory Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody

raised against native TFRC.

Clone Name: MEM-75

Immunogen: Native purified TFRC from NALM-6 human

pre-B cell line.

Host: Mouse

Theoretical MW (kDa): 95

Reactivity: Human

**Applications:** Flow Cyt

(See our web site product page for detailed applications

information)

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

**Specificity:** This antibody reacts with CD71 antigen (transferrin receptor), a 95 KDa type II homodimeric transmembrane glycoprotein expressed on activated B and T lymphocytes, macrophages and erythroid precursors; it is lost on resting blood leukocytes. This antibody does not block binding of transferrin to the receptor.

Form: Liquid

Conjugation: FITC

Isotype: IgG1

**Recommend Usage:** Flow Cytometry (20 ul in human blood cells 100 ul in whole blood or 10<sup>6</sup> cells in a suspension)

The optimal working dilution should be determined by

the end user.

Storage Buffer: In PBS (0.2% BSA, 0.09% sodium

azide)

Storage Instruction: Store in the dark at 4°C. Do not

freeze.

Avoid prolonged exposure to light.

Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 7037

Gene Symbol: TFRC

Gene Alias: CD71, TFR, TFR1, TRFR

## References:

1. Hsa, an adhesin of Streptococcus gordonii DL1, binds to alpha2-3-linked sialic acid on glycophorin A of the erythrocyte membrane. Yajima A, Urano-Tashiro Y, Shimazu K, Takashima E, Takahashi Y, Konishi K. Microbiol Immunol. 2008 Feb;52(2):69-77.

2. Flow cytometric analysis of human bone marrow perfusion cultures: erythroid development and relationship with burst-forming units-erythroid. Rogers CE, Bradley MS, Palsson BO, Koller MR. Exp Hematol. 1996 Apr;24(5):597-604.

3. Cell surface antigen expression in human erythroid progenitors: erythroid and megakaryocytic markers. Nakahata T, Okumura N. Leuk Lymphoma. 1994 May;13(5-6):401-9.