

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

## Datasheet

## Cd19 monoclonal antibody, clone 6D5 (PE/Cy7)

Catalog Number: MAB5694

Regulatory Status: For research use only (RUO)

**Product Description:** Rat monoclonal antibody raised against Cd19.

Clone Name: 6D5

**Immunogen:** Mouse CD19-expressing K562 human erythroleukemia cells.

Host: Rat

Reactivity: Mouse

**Applications:** Flow Cyt, IP (See our web site product page for detailed applications information)

**Protocols:** See our web site at http://www.abnova.com/support/protocols.asp or product page for detailed protocols

**Specificity:** CD19 B-cell differentiation antigen, Mr 95 KDa.

Form: Liquid

Conjugation: PE/Cy7

Isotype: IgG2a, kappa

**Recommend Usage:** Flow Cytometry (0.2 ug/10<sup>6</sup> cells) The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS (0.09% sodium azide)

**Storage Instruction:** Store in the dark at 4°C. Do not freeze. Avoid prolonged exposure to light.

Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 12478

Gene Symbol: Cd19

Gene Alias: AW495831

## References:

 The signaling activity of murine CD19 is regulated during cell development. Krop I, Shaffer AL, Fearon DT, Schlissel MS. J Immunol. 1996 Jul 1;157(1):48-56.
Self-renewal of B-1 lymphocytes is dependent on CD10. Krop L do Equatorilla. AB, Lister DB, Allison M.

CD19. Krop I, de Fougerolles AR, Hardy RR, Allison M, Schlissel MS, Fearon DT. Eur J Immunol. 1996 Jan;26(1):238-42.

3. The CD19-CR2-TAPA-1 complex, CD45 and signaling by the antigen receptor of B lymphocytes. Fearon DT. Curr Opin Immunol. 1993 Jun;5(3):341-8.