

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

## **Datasheet**

## Kit monoclonal antibody, clone 2B8 (Cy5)

Catalog Number: MAB5965

Regulatory Status: For research use only (RUO)

Product Description: Rat monoclonal antibody raised

against native Kit.

Clone Name: 2B8

Immunogen: Native purified Kit from murine bone

marrow mast cells.

Host: Rat

Reactivity: Mouse

**Applications:** Flow Cyt

(See our web site product page for detailed applications

information)

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Specificity: Specificity CD117/c-Kit

Form: Liquid

Conjugation: Cy5

Isotype: IgG2b, kappa

**Recommend Usage:** Flow Cytometry (0.2 ug/10<sup>6</sup> cells) The optimal working dilution should be determined by

the end user.

Storage Buffer: In PBS (0.09% sodium azide)

Storage Instruction: Store in the dark at 4°C. Do not

freeze.

Avoid prolonged exposure to light.

Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 16590

Gene Symbol: Kit

Gene Alias: Bs, CD117, Fdc, Gsfsco1, Gsfsco5, Gsfsow3, SCO1, SCO5, SOW3, Ssm, Tr-kit, W, c-KIT

## References:

- 1. Interaction of stem cell factor and its receptor c-kit mediates lodgment and acute expansion of hematopoietic cells in the murine spleen. Broudy VC, Lin NL, Priestley GV, Nocka K, Wolf NS. Blood. 1996 Jul 1:88(1):75-81.
- 2. Distinct stages of melanocyte differentiation revealed by anlaysis of nonuniform pigmentation patterns. Yoshida H, Kunisada T, Kusakabe M, Nishikawa S, Nishikawa SI. Development. 1996 Apr;122(4):1207-14.

  3. Functional hierarchy of c-kit and c-fms in intramarrow production of CFU-M. Sudo T, Nishikawa S, Ogawa M, Kataoka H, Ohno N, Izawa A, Hayashi S, Nishikawa S. Oncogene. 1995 Dec 21;11(12):2469-76.