

9F, No. 108, Jhouzih St., Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

## **Datasheet**

## Itga4 monoclonal antibody, clone PS/2 (FITC)

Catalog Number: MAB6097

Regulatory Status: For research use only (RUO)

Product Description: Rat monoclonal antibody raised

against native Itga4.

Clone Name: PS/2

Immunogen: Native purified from P815 mastocytoma

cells.

Host: Rat

Reactivity: Mouse

Applications: Flow Cyt, IHC

(See our web site product page for detailed applications

information)

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

**Specificity:** Specificity CD49d, the alpha4 chain (100 KDa) of the VLA-4 integrin heterodimer (Mr 150-KDa)

Form: Liquid

Conjugation: FITC

Isotype: IgG2b, kappa

**Recommend Usage:** Flow Cytometry (2 ug/10<sup>6</sup> cells) The optimal working dilution should be determined by

the end user.

Storage Buffer: In PBS (0.09% sodium azide)

Storage Instruction: Store in the dark at 4°C. Do not

freeze.

Avoid prolonged exposure to light.

Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 16401

Gene Symbol: Itga4

Gene Alias: CD49D

## References:

- 1. Alpha 4 integrin directs virus-activated CD8+ T cells to sites of infection. Christensen JP, Andersson EC, Scheynius A, Marker O, Thomsen AR. J Immunol. 1995 May 15;154(10):5293-301.
- 2. Expression of murine VCAM-1 in vitro and in different models of inflammation in vivo: correlation with immigration of monocytes. Henseleit U, Steinbrink K, Sunderkotter C, Goebeler M, Roth J, Sorg C. Exp Dermatol. 1994 Dec;3(6):249-56.
- 3. Decreased adhesion to endothelial cells and matrix proteins of H-2Kb gene transfected tumour cells. Lauri D, De Giovanni C, Biondelli T, Lalli E, Landuzzi L, Facchini A, Nicoletti G, Nanni P, Dejana E, Lollini PL. Br J Cancer. 1993 Nov;68(5):862-7.