

Datasheet

Rabbit Anti-Bovine IgG F(c) secondary antibody (Rhodamine)

microfluorometric and fluorescence microscopic studies.
Titus JA, Haugland R, Sharrow SO, Segal DM. J
Immunol Methods. 1982;50(2):193-204.

Catalog Number: PAB10458

Regulatory Status: For research use only (RUO)

Product Description: Rabbit anti-bovine IgG recognizes bovine IgG F(c) fragment. This secondary antibody was purified using antigen affinity chromatography. The antibody is conjugated with Rhodamine.

Immunogen: Bovine IgG F(c) fragment

Host: Rabbit

Reactivity: Bovine

Applications: Flow Cyt, IF
(See our web site product page for detailed applications information)

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Lyophilized

Conjugation: Rhodamine (TRITC)

Purification: This product was prepared from monospecific antiserum by immunoaffinity chromatography, followed by solid phase adsorption(s) to remove any unwanted reactivities.

Recommend Usage: The optimal working dilution should be determined by the end user.

Storage Buffer: Lyophilized from 20 mM KH_2PO_4 , 150 mM NaCl, pH 7.2

Storage Instruction: Store at 4°C on dry atmosphere. After reconstitution with 1 mL of deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

References:

1. Texas Red, a hydrophilic, red-emitting fluorophore for use with fluorescein in dual parameter flow