

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

Datasheet

Rabbit Anti-Goat IgG F(c) secondary antibody (Fluorescein)

Catalog Number: PAB10587

Regulatory Status: For research use only (RUO)

Product Description: Rabbit anti-goat IgG recognizes goat IgG F(c) fragment. This secondary antibody was purified using antigen affinity chromatography. The antibody is conjugated with Fluorescein.

Immunogen: Goat IgG F(c) fragment

Host: Rabbit

Reactivity: Goat

Applications: Flow Cyt, IF

(See our web site product page for detailed applications

information)

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Form: Lyophilized

Conjugation: FITC

Purification: This product was prepared from monospecific antiserum by immunoaffinity chromatography, followed by solid phase adsorption(s) to remove any unwanted reactivities.

Conjugation Note: 3.1 moles FITC per mole of rabbit

IgG

Recommend Usage: The optimal working dilution

should be determined by the end user.

Storage Buffer: Lyophilized from 10 mM Sodium Phosphate, 150 mM Sodium Chloride, pH 7.2 (10 mg/mL

BSA, 0.01% thimerosal)

Storage Instruction: Store at 4°C on dry atmosphere. After reconstitution with deionized water, store at -20°C

or lower.

Aliquot to avoid repeated freezing and thawing.

References:

1. Immunofluorescence using dichlorotriazinylaminofluorescein (DTAF). I. Preparation and fractionation of labelled IgG. Blakeslee D, Baines MG. J Immunol Methods. 1976;13(3-4):305-20.