

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

## **Datasheet**

## MYC polyclonal antibody

Catalog Number: PAB26624

Regulatory Status: For research use only (RUO)

Product Description: Rabbit polyclonal antibody raised

against synthetic peptide of MYC.

Immunogen: A synthetic peptide corresponding to

residues surrounding S373 of human MYC.

Sequence: K-R-Sp-F-F

Host: Rabbit

Reactivity: Human, Mouse, Rat

**Applications: WB-Ce** 

(See our web site product page for detailed applications

information)

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Form: Liquid

Purification: Affinity chromatography

Concentration: 1 mg/mL

**Recommend Usage:** Western Blot (1:500-1:1000) The optimal working dilution should be determined by

. . . .

the end user.

**Storage Buffer:** In PBS (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), 150 mM NaCl, pH 7.4 (50% glycerol, 0.02% sodium azide)

Storage Instruction: Store at -20°C.

Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 4609

Gene Symbol: MYC

Gene Alias: bHLHe39, c-Myc

**Gene Summary:** The protein encoded by this gene is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular

transformation. It functions as a transcription factor that regulates transcription of specific target genes. Mutations, overexpression, rearrangement and translocation of this gene have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt lymphoma. There is evidence to show that alternative translation initiations from an upstream, in-frame non-AUG (CUG) and a downstream AUG start site result in the production of two isoforms with distinct N-termini. The synthesis of non-AUG initiated protein is suppressed in Burkitt's lymphomas, suggesting its importance in the normal function of this gene. [provided by RefSeq]