

**Anti-MBL (human)**  
**Mouse monoclonal antibody**  
**HYB 131-10**

Subclass: IgG1/k  
Clone:1E2

CAT. NO.

---

SPECIFICITY	HYB 131-10 is specific for MBL (mannan-binding lectin) from human serum or plasma.
IMMUNOGEN	MBL purified from human donor plasma.
TESTED APPLICATIONS	ELISA, WB
SPECIES REACTIVITY (POSITIVE)	Human
SPECIES REACTIVITY (NEGATIVE)	Not determined
EPITOPE SPECIFICITY	The epitope is thought to be on the carbohydrate recognition domain and differs from that of HYB 131-01 and HYB 131-11.

---

**PRESENTATION**

Content:	Available in 200 µL and 1 mL size. 1 mg/mL +/- 15%. See Certificate of Analysis for details.
Preparation:	Protein-A purified
Form:	Liquid
Solvent:	0.01 M phosphate buffer, pH 7.4, containing 0.5 M NaCl and 15 mM sodium azide
Storage:	4-8°C without exposure to light. No precautions necessary during handling.

---

**APPLICATION**

**ELISA:** HYB 131-10 reacts strongly with MBL. Strong reaction is seen in ELISA with MBL coated directly onto the microtiter well. In Western blotting HYB 131-10 reacts with human MBL in both its oligomerized state and as single protein chain of 26 kDa. HYB 131-10 can be used to measure MBL in plasma or serum but is not specific for oligomerized MBL when used in combination with itself or other MBL antibodies(1).  
**WB:** HYB 131-10 can be used in Western blotting.

**TARGET**

Mannan-binding lectin (MBL), also called mannanose-binding lectin or protein, belongs to the C-type family of collectins, showing calcium-dependent binding to certain sugars. It consists of oligomers of triple-chain subunits and its binding and complement activating activities depend on its normal oligomerization. On binding to mannan-like microbial surface carbohydrates, MBL activates the complement system by means of its own lectin pathway, depending on the MBL-associated serine proteases (MASPs). Because of the presence of different structural and promoter alleles in the population, 12% or more of the population have low concentrations (<50ng/mL) of normally oligomerized, functional MBL in plasma or serum.

**REFERENCES**

1. Hein E, Honoré C, Skjoedt MO, Munthe-Fog L, Hummelshøj T, Garred P. (2010) Functional analysis of Ficolin-3 mediated complement activation. PlosOne 5(11):e15443. doi: 10.1371/journal.pone.0015443.

**CONDITIONS**

Unless otherwise marked, all products are for research use only. Not for use in diagnostic procedures. Not for use in human therapeutic applications. For in vitro use or further manufacture only. The information and product are offered without guarantee as the ultimate conditions of use are beyond our control. The foregoing is in lieu of all warranties, expressed or implied, including implied warranties of merchantability and fitness for a particular purpose. In no event shall BioPorto Diagnostics A/S be responsible for loss of profits or indirect consequential losses resulting from use of its product