

## Acris Antibodies, Inc.

6815 Flanders Drive, Suite 140 San Diego, CA 92121 UNITED STATES Phone: +1-858-888-7900

Phone: +1-858-888-7900 Fax: +1-858-888-7904 <u>US-info@acris-antibodies.com</u>

## Acris Antibodies GmbH

Schillerstr. 5 32052 Herford GERMANY

AM00092PII-N

Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com

## Monoclonal Antibody to MAPKK 2 (incl. pos. control) - Purified

Alternate names: Dual specificity mitogen-activated protein kinase kinase 2, ERK activator kinase 2, MAP

kinase kinase 2, MAP kinase kinase 2, MAPK/ERK kinase 2, MEK2, MKK2

Catalog No.: AM00092PU-N

Quantity: 0.1 mg

Background: MEK (MAP Kinase Kinase) phosphorylates the MAP Kinase on both threonine and tyrosine

residues of the activation motif TEY. MEK1 and MEK2 are activated by phosphorylation of

two serine residues (Ser 218/222 in MEK1 and Ser 222/226 in MEK2). These

phosphorylation sites are substrates of the Raf family of kinases.

Uniprot ID: P36507

NCBI: <u>NP 109587.1</u>

GenelD: <u>5605</u>

Host / Isotype: Mouse / IgG1

Clone: 8E8

Immunogen: Synthetic peptide conjugated to KLH

Format: State: Lyophilized purified IgG

**Purification:** Size exclusion chromatography

Buffer System: 1 ml PBS / 0.09 % Na-azide / PEG and Sucrose

**Reconstitution:** Restore with 1 ml H2O (15 min, RT)

**Applications:** Western Blot: 0.5 μg/ml for HRPO/ECL detection.

Recommended blocking buffer: Casein/Tween 20 based blocking and blot incubation

buffer.

Positive Control: Cell lysate from untreated HepG2 cells.

ELISA: 0.1 µg/ml (protein ELISA).

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

**Specificity:** This antibody specifically recognizes the N-terminus of MEK2 at 45 kDa.

Species: Human, Mouse, Rat, Dog.

Other species not tested.


Storage: Store lyophilized (preferably in a desiccator) at -20°C and reconstituted (aliquote and

freeze in liquid nitrogen) at -80°C. Avoid repeated freezing and thawing.

Thaw aliquots at 37°C. Thawed aliquots may be stored at 4°C up to 3 months.

Shelf life: one year from despatch.

Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com





**Pictures:** 

Detection of endogenous MEK2 Whole cell lysates of serum starved tumor cells (20.000 cells per lane) were applied to SDS-PAGE and transferred to a PVDF membrane. The immunoblot was probed with mab MEK2-8E8(0.5  $\mu$ g/ ml) for 1h at RT and developed by ECL (exp. time: 30 sec). lane 1: A431; lane 2: A549; lane 3: SKOV3; lane 4: OVCAR5; lane 5: HaCaT; lane 6: PC3; lane 7: HeLa; lane 8: HepG2

