

Acris Antibodies, Inc.

6815 Flanders Drive, Suite 140 San Diego, CA 92121 UNITED STATES Phone: +1-858-888-7900

Phone: +1-858-888-7900 Fax: +1-858-888-7904 <u>US-info@acris-antibodies.com</u>

Acris Antibodies GmbH

Schillerstr. 5 32052 Herford GERMANY

AM01316FC-N

Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com

Monoclonal Antibody to EMR3 - FITC

Alternate names: EGF-like module receptor 3, EGF-like module-containing mucin-like hormone receptor-like 3

Catalog No.: AM01316FC-N

Quantity: 0.1 mg
Concentration: 0.1 mg/ml

Background: EMR3 is expressed at the cell surface as a heterodimer. The molecule is predominantly

expressed on granulocytes, and at lower levels on mature myeloid cells, monocytes and dendritic cells. EMR3 is absent on lymphocytes, haematopoietic stem cells and myeloid progenitors. Studies suggest that the EMR3 molecule is up-regulated during late stages of neutrophil differentiation and is a marker for terminally differentiated cells. The exact

functions of EMR3 and its ligands have not yet been determined.

Uniprot ID: Q9BY15

NCBI: NP_115960.2

GenelD: 84658

Host / Isotype: Hamster / IgG

Clone: 3D7

OG/20130615

Immunogen: ARHO-EMR3-CD97 (EGF1) transfectants. Spleen cells from immunised Armenian hamster

were fused with cells of the mouse SP2/0 myeloma cell line.

Format: State: Liquid purified IgG

Purification: Affinity chromatography on Protein G

Buffer System: PBS, pH 7.4, containing 0.09% Sodium Azide

Label: FITC - Fluorescein Isothiocyanate Isomer I

Applications: Flow cytometry (Neat - 1/10): use 10 μl of the suggested working dilution to label 10e6 cells

in 100 μl.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This antibody recognises epidermal growth factor (EGF) module-containing mucin-like

hormone receptor 3 (EMR3), which is a 56kD member of the EGF-7 transmembrane (TM7)

family of adhesion receptors.

Species: Human.

Other species not tested.

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

This product is photosensitive and should be protected from light.

Shelf life: one year from despatch.



AM01316FC-N: Monoclonal Antibody to EMR3 - FITC

General Readings:

1. Stacey M, Lin HH, Hilyard KL, Gordon S, McKnight AJ. Human epidermal growth factor (EGF) module-containing mucin-like hormone receptor 3 is a new member of the EGF-TM7 family that recognizes a ligand on human macrophages and activated neutrophils. J Biol Chem. 2001 Jun 1;276(22):18863-70. Epub 2001 Mar 14. PubMed PMID: 11279179.

