

Acris Antibodies, Inc.

6815 Flanders Drive, Suite 140 San Diego, CA 92121 UNITED STATES

Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com

Acris Antibodies GmbH

Schillerstr. 5 32052 Herford GERMANY

AM05208FC-N

Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com

Monoclonal Antibody to CD11b - FITC

Alternate names: CD11 antigen-like family member B, CR-3 alpha chain, CR3A, Cell surface glycoprotein

MAC-1 subunit alpha, ITGAM, Integrin alpha-M, Leukocyte adhesion receptor MO1, MAC1,

Neutrophil adherence receptor

Catalog No.: AM05208FC-N
Quantity: 100 Tests
Concentration: Lot specific

Background: Mac-1), CD11b/CD18 complex is a leukocyte adhesion receptor to intercellular adhesion

molecules (ICAM's). CR3 alpha chain (M.W. 165,000) is on virtually all peripheral blood neutrophils, eosinophils, monocytes and approximately 30% of peripheral blood

lymphocytes.

Uniprot ID: P11215

NCBI: NP 000623.2

GenelD: <u>3684</u>

Host / Isotype: Mouse / IgG2a

Clone: F6.2

Format: State: Liquid purified IgG fraction.

Buffer System: PBS containing 0.2% protein carrier and 0.08% Sodium Azide as

preservative. **Label:** FITC

Applications: Flow Cytometry.

Monitoring of subsets in peripheral blood. Characterization of leukemias and lymphomas.

Analysis of CTL-mediated cytotoxicity. Studies of transplantation rejection.

Functional studies of cell adhesion molecules.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: Species: Human.

Other species not tested.

Storage: Store the antibody undiluted at 2-8°C.

Do Not Freeze!

Shelf life: One year from despatch.

General Readings: 1. Arnaout MA. Structure and function of the leukocyte adhesion molecules CD11/CD18.

Blood. 1990 Mar 1;75(5):1037-50. PubMed PMID: 1968349.

2. Leukocyte adhesion to endothelial cells. Wautier, ,J.L., Sediadi, H., Vilette, d., Weill, D.,

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service

Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com





AM05208FC-N: Monoclonal Antibody to CD11b - FITC

Wautier, M.P.. Biorheology . 1990:27(3-4): 425-32.

- 3. Babcock GF, Alexander JW, Warden GD. Flow cytometric analysis of neutrophil subsets in thermally injured patients developing infection. Clin Immunol Immunopathol. 1990 Jan;54(1):117-25. PubMed PMID: 1967160.
- 4. Leukocyte Adhesion Deficiency: an inherited defect in Mac-1,LFA-1, and p150,95 glycoproteins. Anderson, D.C. and Springer, T.A.. Ann. Rev. Med. 1987;38:175.
- 5. Merrill JT, Slade SG, Weissmann G, Winchester R, Buyon JP. Two pathways of CD11b/CD18-mediated neutrophil aggregation with different involvement of protein kinase C-dependent phosphorylation. J Immunol. 1990 Oct 15;145(8):2608-15. PubMed PMID: 1976698.
- 6. Richter J, Ng-Sikorski J, Olsson I, Andersson T. Tumor necrosis factor-induced degranulation in adherent human neutrophils is dependent on CD11b/CD18-integrintriggered oscillations of cytosolic free Ca2+. Proc Natl Acad Sci U S A. 1990 Dec;87(23):9472-6. PubMed PMID: 1979172.

Protocols:

PBMC:

Add 10 ul of MAB/10e6 PBMC in 100 ul PBS.

Mix gently and incubate for 15 minutes at 2-8°C.

Wash twice with PBS and analyze or fix with $0.5\% \, v/v$ of paraformaldehyde in PBS and analyze.

WHOLE BLOOD:

Add 10 ul of MAB/100 ul of whole blood.

Mix gently and incubate for 15 minutes at room temperature (20 $^{\circ}\text{C}).$

Lyse the whole blood.

Wash once with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze.

See instrument manufacturers instructions for Lysed Whole Blood and Immunofluorescence analysis with a flow cytometer or microscope.



Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com