

## Acris Antibodies, Inc.

6815 Flanders Drive, Suite 140 San Diego, CA 92121 UNITED STATES Phone: +1.858.888.7900

Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com AM05883FC-N

## **Acris Antibodies GmbH**

Schillerstr. 5 32052 Herford GERMANY

Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com

## Monoclonal Antibody to CD11R3 - FITC

Catalog No.: AM05883FC-N

Quantity: 0.1 mg
Concentration: 0.1 mg/ml
Host / Isotype: Mouse / IgG1
Clone: 2F4/11

Immunogen: Porcine alveolar macrophages

Format: State: Liquid Ig fraction

Purification: Affinity chromatography on Protein G

Buffer System: Phosphate buffered saline pH7.4, 0.09% Sodium Azide, 1% Bovine Serum

Albumin

Label: FITC - Fluorescein Isothiocyanate Isomer 1

Applications: Flow Cytometry: 10 μl of neat - 1:10 diluted antibody to label 1x10e6 cells in 100 μl.

Functional assays: Removal of sodium azide is recommended prior to use.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This antibody recognises porcine CD11R3, a 155kD cell surface marker which is a member of

the alpha integrin family. CD11R3 has a similar expression pattern to the human CD11b marker, being expressed on granulocytes, monocytes and alveolar macrophages, but not on lymphocytes, eythrocytes or platelets. Clone 2F4/11 is reported to block phagocytosis of complement-opsonised zymosan particles by polymononuclear granulocytes and alveolar

macrophages (1). **Species:** Porc.

Other species not tested.

Storage: Store the antibody at 2 - 8 °C up to one month or (in aliquots) at -20 °C for longer. Avoid

repeated freezing and thawing.

This aproduct is photosensitive and should pe protected from light.

Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf life: one year from despatch.

General Readings: 1. Bullido R, Alonso F, Gómez del Moral M, Ezquerra A, Alvarez B, Ortuño, et al. Monoclonal

antibody 2F4/11 recognizes the alpha chain of a porcine beta 2 integrin involved in adhesion and complement mediated phagocytosis. J Immunol Methods. 1996 Sep

9;195(1-2):125-34. PubMed PMID: 8814327.

2. Domínguez J, Alvarez B, Alonso F, Thacker E, Haverson K, McCullough K, et al. Workshop studies on monoclonal antibodies in the myeloid panel with CD11 specificity. Vet Immunol

Immunopathol. 2001 Jul 20;80(1-2):111-9. PubMed PMID: 11445222.

3. Sánchez-Torres C, Gómez-Puertas P, Gómez-del-Moral M, Alonso F, Escribano JM,

Ezquerra A, et al. Expression of porcine CD163 on monocytes/macrophages correlates with

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com

Antibody Hotline - Technical Questions - Antibody Location Service





## AM05883FC-N: Monoclonal Antibody to CD11R3 - FITC

permissiveness to African swine fever infection. Arch Virol. 2003 Dec;148(12):2307-23. Epub 2003 Sep 16. PubMed PMID: 14648288.

- 4. Van de Walle GR, Favoreel HW, Nauwynck HJ, Mettenleiter TC, Pensaert MB. Transmission of pseudorabies virus from immune-masked blood monocytes to endothelial cells. J Gen Virol. 2003 Mar;84(Pt 3):629-37. PubMed PMID: 12604815.
- 5. Alvarez B, Doménech N, Alonso F, Sánchez C, Gómez del Moral M, Ezquerra A, et al. Molecular and functional characterization of porcine LFA-1 using monoclonal antibodies to CD11a and CD18. Xenotransplantation. 2000 Nov;7(4):258-66. PubMed PMID: 11081760.

