

Monoclonal Antibody to CD16/CD32 - FITC

Alternate names:	CD16a, FCG3, FCGR3A, Fc gamma receptor IIB, Fc-gamma RIII, Fc-gamma RIII-alpha, Fc-gamma RIIIa, Fc-gamma-RIIB, FcR mouse Seroblock, FcRII, FcRIII, FcRIIIa, Fcgr2, IGF3, IgG Fc receptor II beta, IgG Fc receptor III-2, Low affinity immunoglobulin gamma Fc region receptor II/III, Ly-17, Lymphocyte antigen 17
Catalog No.:	AM08040FC-N
Quantity:	0.5 mg
Concentration:	0.5 mg/ml
Background:	<p>The lymphocyte Fc-gamma Receptors recognize the Fc portion of IgG, presented either as immune complexes or as free antibody. The different classes of receptors are distinct because of varying size, tissue distribution and affinity for IgGs. The Fc type II receptor is expressed on a wide variety of cells including B cells, hematopoietic cells, monocyte/macrophages, neutrophils, platelets, Langerhans cells, eosinophils, basophils, trophoblasts, and endothelial cells of the placenta. (Ref.1,2)</p> <p>The Fc-gamma type III receptors are higher affinity than the type II and are expressed on macrophages, NK cells and neutrophils. Both types of receptors can be expressed on the same cell and in varying ratios. (Ref.1) The receptors are constitutively expressed, although cytokines and lymphokines can modulate their expression. (Ref.4) Besides identifying Fc-gammaR+ cells, monoclonal antibodies to the Fc-gammaII/III receptor have been used to block Fc receptor binding of IgG, Fc-mediated signal transduction and effector functions, clearance of immune complexes and to attenuate infection by organisms dependent on Fc-gamma R for parasitic invasion. (Ref.4-6)</p>
Host / Isotype:	Rat / IgG2a
Clone:	93
Format:	State: Liquid purified Ig fraction. Buffer System: PBS containing 0.09% Sodium Azide as preservative. Label: FITC – Fluorescein Isothiocyanate Isomer 1
Applications:	Flow Cytometry: < / 3 µg/10e6 cells. (Ref.4,7,8) Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody is specific to Murine CD16/32 (CD16/FcγII and CD32/FcγIII receptors). It recognizes a conformational epitope formed by Fc-gammaII and Fc-gamma III Receptors. Species: Mouse. Other species not tested.
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing. Shelf life: one year from despatch.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com



- General Readings:**
1. Unkeless, J. C., et al. 1995. Semin. Immunol. 7:37.
 2. Warmerdam, P.A., et al. 1992. Immunobiol. 185:175.
 3. Huizinga, T. W., et al. 1989. J. Immunol. 142:2365.
 4. Ho, A. S-Y., et al. 1995. Mol. and Cell. Biol. 15:5043.
 5. John Kearney, personal communication.
 6. Kurlander, R.J., D. M. Ellison and J. Hall. 1984. J. Immunol. 133:855.
 7. Araujo-Jorge, T., et al. 1993. Infection and Immunity 61:4925.
 8. Titus, J. A., et al. 1984. J. Immunol. 133:556.

Pictures:

Immunofluorescent Staining:
BALB/c spleen cells were double-stained with Rat anti-Mouse CD16/32-FITC and Rat anti-Mouse CD19-R-PE (clone 6D5). Lymphocytes were gated and analyzed on a FACScan(TM) flow cytometer (BDIS, San Jose, CA). Amount Used: 3 µg/10e6 cells.

