

Monoclonal Antibody to KLRG1 - PE

C-type lectin domain family 15 member A, CLEC15A, ITIM-containing receptor MAFA-L, Alternate names:

KLRG1, Killer cell lectin-like receptor subfamily G member 1, MAFA, MAFA-like receptor,

MAFAL, Mast cell function-associated antigen

AM08067RP-S **Catalog No.:**

Quantity: 0.1 mg **Concentration:** $0.1 \, \text{mg/ml}$

Background: Killer cell lectin-like receptor G1 (KLRG1; formerly known as mouse MAFA or 2F1 Ag), a

> homodimeric member of the lectin-like type 2 transmembrane receptor family that contain characteristic immunoreceptor tyrosine-based inhibitory motifs (ITIMs) in their cytoplasmic domains. (Ref.1,2) These ITIMs interact with the SH2 domains of protein phosphatases such as SHP1. (Ref.3,4) Cell surface expression of KLRG1 is up-regulated by expression of MHC class I molecules. The effect of MHC class I expression is indirect, and can be mediated by interactions with class I-specific Ly49 inhibitory receptors. (Ref.1)

Uniprot ID: 088713

NCBI: NP 058666.1

GeneID: 50928

Host / Isotype: Hamster / IgG

Clone: 2F1

Immunogen: Activated NK (A-LAK) cells from B6 (H-2b) mice. (Ref.1)

Format: State: Liquid purified Ig fraction.

Buffer System: PBS containing 0.09% Sodium Azide as preservative and a stabilizing

Label: PE - R-Phycoerythrin

Applications: Flow Cytometry: $\langle / = 0.2 \,\mu g/10e6 \, cells. \, (Ref. 1,5)$

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

This antibody reacts with Killer Cell Lectin-like Receptor G1 (KLRG1 formerly known as **Specificity:**

mouse MAFA or 2F1-Ag).

It stains 30-60% of NK1.1+CD3- splenocytes, and a small fraction of T cells in all mouse

strains tested (C57BL/6, BALB/c, 129/J, C3H.SW, AKR/J, SJL). (Ref.1,5)

Species: Mouse.

Other species not tested.





AM08067RP-S: Monoclonal Antibody to KLRG1 - PE

Storage: Store the antibody undiluted at 2-8°C.

DO NOT FREEZE!

This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing. Shelf life: one year from despatch.

General Readings:

1. Corral, L., T. Hanke, R.E. Vance, D. Cado, and D.H. Raulet. 2000. Eur. J. Immunol. 30:920.

2. Hanke, T., L. Corral, R.E., Vance, and D.H. Raulet. 1998. 28:4409.

3. Burshtyn, D., A. Scharenberg, N. Wagtmann, S. Rajagopalan, M. Peruzzi, J.-P. Kinet, and

E.O. Long. 1997. Immunity 4:77.

4. Nakamura, M.C., E.C. Niemi, M.J. Fisher, L.D. Schultz, W.E. Seaman, and J.C. Ryan. 1997.

J. Exp. Med. 185:673.

5. Liu, S., and D.H. Raulet. Personal communication.

